

Deep Carbon Emissions from Volcanoes

Michael R. Burton

*Istituto Nazionale di Geofisica e Vulcanologia
Via della Faggiola, 32
56123 Pisa, Italy
burton@pi.ingv.it*

Georgina M. Sawyer

*Laboratoire Magmas et Volcans, Université Blaise Pascal
5 rue Kessler, 63038 Clermont Ferrand, France
and
Istituto Nazionale di Geofisica e Vulcanologia
Via della Faggiola, 32
56123 Pisa, Italy*

Domenico Granieri

*Istituto Nazionale di Geofisica e Vulcanologia
Via della Faggiola, 32
56123 Pisa, Italy*

INTRODUCTION: VOLCANIC CO₂ EMISSIONS IN THE GEOLOGICAL CARBON CYCLE

Over long periods of time (~Ma), we may consider the oceans, atmosphere and biosphere as a single exospheric reservoir for CO₂. The geological carbon cycle describes the inputs to this exosphere from mantle degassing, metamorphism of subducted carbonates and outputs from weathering of aluminosilicate rocks (Walker et al. 1981). A feedback mechanism relates the weathering rate with the amount of CO₂ in the atmosphere via the greenhouse effect (e.g., Wang et al. 1976). An increase in atmospheric CO₂ concentrations induces higher temperatures, leading to higher rates of weathering, which draw down atmospheric CO₂ concentrations (Berner 1991). Atmospheric CO₂ concentrations are therefore stabilized over long timescales by this feedback mechanism (Zeebe and Caldeira 2008). This process may have played a role (Feulner et al. 2012) in stabilizing temperatures on Earth while solar radiation steadily increased due to stellar evolution (Bahcall et al. 2001). In this context the role of CO₂ degassing from the Earth is clearly fundamental to the stability of the climate, and therefore to life on Earth. Notwithstanding this importance, the flux of CO₂ from the Earth is poorly constrained. The uncertainty in our knowledge of this critical input into the geological carbon cycle led Berner and Lagasa (1989) to state that it is the most vexing problem facing us in understanding that cycle.

Notwithstanding the uncertainties in our understanding of CO₂ degassing from Earth, it is clear that these natural emissions were recently dwarfed by anthropogenic emissions, which have rapidly increased since industrialization began on a large scale in the 18th century, leading to a rapid increase in atmospheric CO₂ concentrations. While atmospheric CO₂ concentrations have varied between 190-280 ppm for the last 400,000 years (Zeebe and Caldeira 2008), human activity has produced a remarkable increase in CO₂ abundance, particularly in the last 100 years, with concentrations reaching ~390 ppmv at the time of writing. This situation highlights

the importance of understanding the natural carbon cycle, so that we may better determine the evolution of the anthropogenic perturbation.

The principle elements of the multifaceted and complex geological carbon cycle are summarized in Figure 1. The main sources of carbon are active and inactive volcanism from arcs and rift zones and metamorphism of crustal carbonates. The main sinks for geological carbon are silicate weathering and carbonation of oceanic crust. Knowledge of both the total magnitude of carbon ingassing during subduction and carbon released from volcanism and metamorphism would allow quantification of the evolution and relative distribution of volatiles in the crust and mantle (Dasgupta and Hirschmann 2010).

The main focus of this work is the role of volcanism in producing CO₂ in the atmosphere and oceans. Volcanic CO₂ sources can be divided into several categories, direct and diffuse degassing from active arc and rift volcanoes, diffuse degassing from inactive volcanoes and regional diffuse degassing from intrusive plutonic structures with associated crustal metamorphism. We focus here on non-eruptive degassing because, as shown below, continuous emission of CO₂ from multiple sources appears to dominate short-lived eruptive emissions from point sources.

CO₂ released directly from active volcanoes has three main sources, CO₂ dissolved in the mantle, recycled CO₂ from subducted crustal material (e.g., Marty and Tolstikhin 1998) and decarbonation of shallow crustal material (e.g., Troll et al. 2012). Separating the relative proportions of mantle and crustal carbon is possible through investigation of the isotopic composition of emitted carbon (e.g., Chiodini et al. 2011) and is increasingly important given that during eruptions magmatic intrusions may interact with crustal material, strongly enhancing the CO₂ output of the volcanic system (Troll et al. 2012), at least temporarily. The magnitude of diffuse mantle CO₂ can also be identified isotopically in mixed metamorphic and magmatic gases using Carbon (Chiodini et al. 2011) or Helium isotopes as a proxy for deep mantle sources in both major fault systems (Pili et al. 2011) and crustal tectonic structures (Crossey et al. 2009).

Our current estimates of volcanic carbon emissions are poorly constrained due to a lack of direct measurements. Measuring CO₂ in subaerial volcanic plumes is a challenge, because while CO₂ typically makes up ~10 mol% of volcanic gas emissions (the majority of which is normally water vapor), mixing with the atmosphere rapidly dilutes the volcanic CO₂ signature. Nevertheless, technological advances and an increase in the number of volcanoes studied have




Figure 1. Schematic diagram showing the main sources and sinks for volcanic CO₂ on Earth within the geological carbon cycle. CO₂ is released at MORs during crustal genesis, but it is also absorbed into the newly formed crust in high temperature hydrothermal reactions. Carbonates precipitate directly into oceanic crust and collect in sediments before the subducting slab is carried under the mantle wedge. Volcanism then produces CO₂ emissions from the fore-arc (through cold seeps), arc volcanism, rift volcanism, intra-continental volcanoes and hotspots. The CO₂ emitted into the atmosphere reacts in weathering reactions with silicate rocks, carrying C back into the ocean where the geological carbon cycle is eventually closed through carbon sequestration into the subducting slab material.

greatly increased our knowledge of volcanic CO₂ fluxes over the last 10 years. One of the main goals of this work is to update global geological CO₂ flux estimates (e.g., Kerrick 2001; Mörner and Etiope 2002; Fischer 2008) using the recently acquired volcanic CO₂ flux data.

Diffuse CO₂ degassing from both volcanic and tectonic structures is a large contributor to the global geological CO₂ emission, but is difficult to measure due to the large areal extent that may be in play, and the large number of degassing sites throughout the globe. Measuring the CO₂ degassing rates into volcanic lakes and from submarine volcanism have significant technical challenges. In the following we review the state of the art of volcanic CO₂ measurements and present a catalogue of reported, quantified emissions from geological sources. These measurements are then extrapolated to produce estimates of the global volcanic CO₂ flux. These estimates are compared with previously published estimates of total CO₂ emissions, silicate weathering rates and the rate of carbon consumption during subduction. We then examine the dynamic role of CO₂ within magmatic systems and the magnitude of CO₂ released during eruptions.

Carbon species in Earth degassing

CO₂ is not the only carbon-containing molecule emitted from the Earth. In order of decreasing emissions, CO₂, CH₄, CO and OCS all contribute to the total carbon budget. Mörner and Etiope (2002) estimated that the global emission of CO₂ from Earth degassing was ~600 million tonnes of CO₂ per year (Mt/yr, 1 Mt = 10¹² g), with ~300 Mt/yr produced from subaerial volcanism, and another 300 Mt/yr produced from non-volcanic inorganic degassing, mostly from tectonically active areas (Chiodini et al. 2005). For comparison, Cadle (1980) estimated that volcanic activity produces 0.34 Mt/yr of CH₄. Mud volcanoes in Azerbaijan were estimated to produce ~1 Mt/yr of CH₄, however the global flux from mud volcanism is not known. Hydrocarbon seepage of CH₄ globally is estimated to produce between 8 and 68 Mt/yr (Hornafius et al. 1999). Etiope et al. (2008) estimated that global CH₄ emissions from geological sources to be 53 Mt/yr, a significant proportion of the geological C output.

CO is emitted directly from volcanoes, with a CO₂/CO ratio that varies between ~10 and ~1000 depending on the oxygen fugacity and temperature of the fluid co-existing with melt prior to outgassing. OCS is also directly emitted within volcanic plumes, but in even smaller relative amounts than CO, typically 1000-10,000:1 for CO₂:OCS (Mori and Notsu 1997; Burton et al. 2007a; Oppenheimer and Kyle 2008; Sawyer et al. 2008a). OCS is the most abundant S bearing gas species in the atmosphere, contributes to stratospheric sulfuric acid aerosol generation (Crutzen 1976) and is an efficient greenhouse gas (Brühl et al. 2012). Its budget is dominated by emissions from oceans and anthropogenic processes. From a total global output of ~1.3 Mt/yr of OCS only 0.03 Mt is estimated to arise from volcanism (Watts 2000). CS₂ is the final trace carbon gas emission from volcanoes, with a similar flux and chemistry to OCS.

While recent estimates of geological emissions of CH₄ (Etiope et al. 2008) clearly indicate that these emissions are significant compared with geological CO₂ on the global scale, in this work we focus on CO₂ emissions, and use the most recent volcanic CO₂ flux data to update the measured global volcanic CO₂ flux.

METHODS FOR MEASURING GEOLOGICAL CO₂ EFFLUX

Ground-based measurements of volcanic plumes

Directly quantifying volcanic CO₂ fluxes in the atmosphere is challenging due to the relatively abundant concentration of background CO₂, currently ~390 ppm. To put this in context, a strong volcanic CO₂ source such as Mt. Etna, Italy, produces a gas plume where 1 km downwind the average concentration of volcanic CO₂ is only ~4 ppm (based on calculations with VOLCALPUFF plume dispersal model, Barsotti et al. 2008). Thus, an *in situ* measure-

ment 1 km downwind needs to resolve a mere 1% excess CO₂ compared with the background concentration. Most volcanoes emit less CO₂ than Mt. Etna, so this is an optimistic scenario. This difficulty has led most researchers to focus on measurements of the volcanic emissions close to the source, using *in situ* and proximal remote sensing techniques. In such measurements, volcanic CO₂ fluxes are normally determined by measuring the ratio of volcanic CO₂ to another volcanic gas, typically SO₂ due to the ease with which its flux can be measured (Oppenheimer et al. 2011), and then calculating the CO₂ flux as the product of the CO₂/SO₂ ratio and the SO₂ flux. The objective of the majority of the following techniques in the context of quantifying CO₂ fluxes is therefore the determination of the CO₂/SO₂ ratio in the volcanic gas.

An exception to this combined CO₂/SO₂ and SO₂ flux approach was adopted by Marty and Le Cloarec (1992) who utilized global volcanic flux estimates of ²¹⁰Po and ²¹⁰Po/CO₂ ratios measured in direct sampling (see below) to derive estimates of global CO₂ fluxes.

Direct sampling of a volcanic gas can be achieved with the use of Giggenbach bottles (Giggenbach and Goguel 1989), where high temperature fumarolic gases are collected in an alkaline solution for later laboratory analysis. This approach allows both bulk and trace gas species to be quantified, but the fact that the most abundant gas component, H₂O, can condense in the tube leading to the alkali solution bottle means that H₂O is challenging to quantify, and therefore absolute concentrations of the other species can be difficult to define. Air contamination is difficult to avoid, and can further increase the difficulty in determining the original volcanic gas concentrations. Such measurements require working in extremely close proximity to the degassing vent, and are ideally performed on the hottest and most highly pressurized emissions (to avoid air contamination), making their collection challenging and potentially hazardous. In addition, some of the most voluminous volcanic gas sources release very little volatiles from fumaroles, instead the bulk of the emission is open-vent degassing from craters. These plume emissions are impossible to sample without air contamination with such an approach.

The MultiGas approach (Shinohara 2005) has greatly simplified the measurement of volcanic CO₂/SO₂ ratios, allowing automatic, unattended analysis of volcanic plumes for extended periods of time (Aiuppa et al. 2007). This instrument combines a near-infrared spectroscopic measurement of CO₂ and H₂O concentrations with a solid-state chemical sensor for quantification of SO₂ and H₂S. The relatively low cost, low power requirement and ease of use of the instrument make it probably the most convenient and cost-effective way of determining *in situ* CO₂/SO₂ ratios available today. Some potential errors can arise, however, due to the different response time of the CO₂ and SO₂ sensors. Typical response times for the near-infrared optical technique used to measure CO₂ is ~1 s, while response times for SO₂ chemical sensors is typically longer, e.g., ~13–31 s for an Alphasense chemical sensor (Roberts et al. 2012). This means that fast changes in chemical composition or concentration are challenging to capture, however through the use of longer integration times problems arising from diverse sensor response times can be avoided. Quickly changing gas concentrations could instead be captured in theory with an optically based SO₂ measurement.

Remote sensing measurements of CO₂ amounts can be performed with infrared spectroscopy if the volcanic gas concentration is sufficiently high above the background atmospheric CO₂ amount. The first such pioneering infrared measurements of volcanic CO₂ amounts were conducted remarkably early, in 1969 by Naughton et al. (1969), during a lava fountain on Kīlauea, Hawai'i. Since then, open-path Fourier transform infrared (OP-FTIR) spectroscopy (Mori and Notsu 1997) has become a well-utilized tool to measure *in situ* volcanic gas compositions. Modern FTIR spectrometers are light (~8 kg), relatively low power (~30 W) and require no cryogenic cooling (e.g., La Spina et al. 2010), yet allow the simultaneous measurement of many volcanic gases, including H₂O, CO₂, SO₂, HCl, HF, CO, OCS and SiF₄. Measurements of volcanic CO₂ require an infrared light source, either an infrared lamp (Burton et al. 2000) or hot volcanic rocks (Allard et al. 2005; Burton et al. 2007a; Sawyer et al. 2008a). Infrared radiation

is absorbed by volcanic gases before being measured with the infrared spectrometer and the resulting spectra can be analyzed (Burton et al. 2000, 2007a) to produce relative abundances of volcanic H₂O, CO₂, SO₂ and several other gases. Smith et al (2011) concluded that such analytical approaches could deliver accuracies of ~5% in CO₂ amounts. The greatest challenge in performing OP-FTIR measurements is obtaining a suitable source of IR radiation, with sufficient volcanic gas between the source and the spectrometer. Recent innovations in applying the remote-controlled mode at the summit of Stromboli (La Spina et al. 2013) show that it can be used for high temporal resolution monitoring of multiple gas sources.

CO₂ emissions from fumarole fields can be determined through the introduction of a known flux of a tracer gas, such as SF₆, and then measuring the volcanic CO₂/SF₆ ratio in the downwind gas emission. Mori et al. (2001) successfully used this approach to measure CO₂ flux emissions from fumarolic vents on Izu-Oshima (Japan), Kirishima (Japan) and Teide (Canary Islands, Spain).

A recent innovation has been the use of portable mass spectrometers to determine *in situ* volcanic gas compositions (Diaz et al. 2010). This approach was used successfully before and after the 5th January 2010 eruption of Turrialba (Costa Rica), revealing significant changes in CO₂, SO₂ and He concentrations.

Volcanic SO₂ flux measurements

As described above the determination of CO₂ flux requires a further step after measurement of CO₂/SO₂ ratios, multiplication with an SO₂ flux. Several assessments of arc volcanic CO₂ emissions have been produced (e.g., Hilton et al. 2002; Fischer and Marty 2005) by interpreting volcanic SO₂ inventories (e.g., Andres and Kasgnoc 1998; Halmer et al. 2002) Since errors on the SO₂ flux propagate into the CO₂ flux we briefly examine here the methods used to measure SO₂ flux, together with their associated errors. The SO₂ flux is much easier to measure directly than the CO₂ flux for two reasons: firstly, SO₂ is not present in the unpolluted troposphere and secondly, SO₂ has a convenient, relatively strong absorption band in the ultraviolet, easily accessible using scattered sunlight as a source. This has allowed the creation of automatic networks of UV scanners, that permit volcanic SO₂ fluxes to be monitored (Edmonds et al. 2003; Burton et al. 2009; Galle et al. 2010; Oppenheimer et al. 2011). Typically in the literature the greatest quoted source of error in SO₂ flux measurements derives from wind speed estimates, and this error is normally indicated to be ~20-30%. A recent innovation in ground-based measurements of SO₂ fluxes is the SO₂ imaging camera, which uses an imaging sensor sensitive to the UV and optical filters to produce specific sensitivity to SO₂ (e.g., Mori and Burton 2006). This approach has the potential to correct implicitly for wind velocity but it suffers potentially from cross-talk between SO₂ and volcanic ash or aerosol.

Recent work on subtle radiative transfer issues relating to ground-based UV SO₂ flux measurements highlight that there could be large, previously ignored, errors associated with light dilution (Kern et al. 2012). This is a process where light scattering from below the volcanic plume enters the instrument, diluting the light which passed through the volcanic plume from above, resulting in a net underestimation in the SO₂ flux, of up to 90%. The true significance and importance of this effect, and the number of published SO₂ fluxes that require re-analysis, has yet to be evaluated. It should be noted therefore that estimates of global CO₂ flux based on published SO₂ flux data may be subject to revision, depending on the impact of light dilution on SO₂ flux measurements.

In addition to ground-based measurements of SO₂ flux, satellite-based measurements are often used, working in both the ultraviolet (e.g., OMI, SCIAMACHY) and infrared (e.g., MODIS, ASTER, IASI) wavelengths. These instruments can produce maps of SO₂ abundance with a repeat time of ~days. With such a repeat rate the same plume is normally not observed in two different images, and it is therefore not possible to perform a simple cross-correlation

to determine plume velocity and therefore flux from images of SO₂ abundance. The conversion from SO₂ abundance image to a quantitative degassing rate is therefore not trivial, because both the age and velocity of the plume at each point in the image must be derived using an independent method (e.g., Merucci et al. 2011). An additional challenge with satellite-based measurements of SO₂ flux is that the sensitivity to SO₂ decreases in the lower atmosphere, such that low-lying volcanoes are difficult to measure unless they are in eruption, and even higher altitude volcanoes require a relatively large SO₂ degassing rate to be reliably quantified (Carn et al. in press).

Measurements of CO₂ flux using the combination of CO₂/SO₂ ratios and SO₂ fluxes therefore reflect uncertainties in both measurements, typically estimated to be 10-20% and 25-30% respectively. As mentioned above, recent work in radiative transfer analysis highlights the potentially important, but largely overlooked, role that light dilution may play in producing potentially significant underestimates of SO₂ fluxes from UV measurements (Kern et al. 2012). Notwithstanding these drawbacks, at the current time these data are the main constraints available for subaerial deep CO₂ output from volcanoes.

Recent technological advances have allowed such integrated CO₂/SO₂, SO₂ flux measurements to be fully automated for the first time, allowing real-time monitoring of CO₂ fluxes. This is particularly important because the low solubility of CO₂ in magmas means that deep, pre-eruptive, magmatic intrusions may be heralded at the surface by increases in CO₂ flux. Aiuppa et al. (2011) reported the first time series of CO₂ flux collected using combined MultiGas and SO₂ flux networks (Burton et al. 2009) on Stromboli volcano. These revealed distinct oscillations in CO₂ emissions, with periods of relatively high CO₂ degassing followed by periods of low CO₂ degassing, producing a steady average CO₂ degassing rate of ~550 t/d. Such a pattern suggests a steady state supply of CO₂ which is modulated by gas accumulation / permeability / magma supply processes within the magma feeding system. Interestingly, more intense explosive activity was observed after a period of intense CO₂ degassing, opening the possibility of using such observations to forecast explosive volcanic activity at this volcano.

Airborne measurements of volcanic plumes

Two main approaches have been used to measure volcanic CO₂ fluxes from the air. A direct method consists of flying an *in situ* CO₂ analyzer (typically a closed-path near-infrared spectrometer) in a raster or ladder traverse across the cross-section of a volcanic plume (Gerlach et al. 1997; Werner et al. 2008). The resulting data can be interpolated to produce a CO₂ concentration map, which can be integrated over the cross-sectional area of the plume and multiplied with wind speed to produce a CO₂ flux. This approach has the advantage of being a direct measurement of the CO₂ emissions, however each measurement requires ~1 hour and therefore very stable wind conditions are required in order to avoid errors in the flux calculation.

The second airborne approach combines plume traverse measurements with an ultraviolet spectrometer to derive SO₂ flux with *in situ* measurement of the CO₂/SO₂ ratio provided with a closed-path FTIR spectrometer, sampling ambient air as the plane flies through the plume (Gerlach et al. 1998). This is probably the most robust methodology currently available for measuring CO₂ fluxes, as the SO₂ flux analysis can be performed close enough to the plume that light dilution is insignificant. Flying has its own challenges, however, due to the technical constraints involved in performing measurements on a vibrating platform, as well as the costs associated with flight time and difficulties presented from flying within the volcanic plume.

Airborne measurements of volcanic CO₂ may also be achieved by viewing infrared radiance from the ground through a volcanic plume with a hyperspectral radiometer, and this has been successfully demonstrated on Kilauea, Hawai'i using the AVIRIS hyperspectral imager (Spinetti et al. 2008). Results obtained with AVIRIS agreed well with ground-based measurements of CO₂ emissions, validating the method.

More recently, similar measurements to those performed by Gerlach et al. (1998) have been conducted from an unmanned aerial vehicle platform (UAV) (McGonigle et al. 2008). Such an approach is appealing due to the significantly reduced cost and risk, as well as increased accessibility. However, the legal framework for conducting such measurements is complex and varies greatly between countries, making general take-up of such methodologies so far quite limited. Future longer distance stand-off measurements may be possible using larger UAVs offering an intermediate option to space-based satellite retrievals of air column soundings.

Space-based measurements of volcanic plumes

Global measurements of volcanic CO₂ emissions would be the ideal approach to quantifying the global volcanic deep CO₂ budget. The most promising observing platform for volcanic CO₂ is the aptly named Orbiting Carbon Observatory (OCO) (Crisp et al. 2004). The rocket carrying the OCO failed to reach orbit when launched in 2009, and a new launch with a replacement satellite is currently planned for 2014. The OCO utilizes a single telescope to feed light to spectrometers which measure the columnar abundance of both O₂ and CO₂, using absorption bands at wavelengths of 0.67 micron for O₂, and 1.61 and 2.06 micron for CO₂. The purpose of the O₂ column amount measurement is to normalize the CO₂ column measurement to an average CO₂ mole fraction in ppmv. The final error on the average CO₂ concentration in the column is 0.3 wt%. The footprint of the OCO will be 1 km by 1.5 km at nadir, and will have a repeat observation period of 16 days.

A simple calculation of the CO₂ emission from a strong emitter such as Mt. Etna allows a direct estimation of the feasibility of such measurements with OCO. Etna degasses CO₂ at an average rate of 16,000 t/d or ~190 kg s⁻¹ (Allard et al. 1991; Aiuppa et al. 2006, 2008; La Spina et al. 2010). Assuming an optimal geometry in which the gas source was at one edge of a single OCO footprint and the entire plume contained within the 1.5 km length of the pixel, with a windspeed of 5 m s⁻¹ the maximum age of CO₂ in the footprint would be ~300 seconds, and the total volcanic CO₂ mass would be ~57,000 tonnes. Converting this CO₂ mass to molecules and averaging over the OCO nadir footprint area produces a vertical column amount of ~5 × 10¹⁹ molecules cm⁻². The atmospheric vertical column amount of CO₂ at the average altitude of Etna is ~6 × 10²¹ molecules cm⁻² and therefore the volcanic signal would be ~0.8 % of the atmospheric column, which is above the 0.3% error limit of OCO. A slower wind would produce a higher volcanic CO₂ amount, while a less optimal geometry would decrease the relative contribution from the volcano. OCO therefore has the potential for measuring passive CO₂ emissions from Mt. Etna, in optimal conditions. During eruptions the CO₂ emission rate would increase, allowing for easier detection. The majority of degassing volcanoes are less productive than Mt. Etna, however, and would present a challenge for detection from OCO, unless they were undergoing an eruption.

Ground-based measurements of diffuse deep CO₂

Significant amounts of diffuse CO₂ are released from active volcanic areas, not only during eruptions but also during quiescent periods. This volcanic CO₂ discharge occurs over the flanks of the volcanic edifice as diffuse soil emanations (Allard et al. 1991; Baubron et al. 1990), and adds to the voluminous and more obvious degassing from fumaroles and summit craters. Many CO₂ soil flux measurement techniques have been applied to quantify these gases and include both direct and indirect methods (e.g., Reiners 1968; Kucera and Kirkham 1971; Kanemasu et al. 1974; Parkinson 1981).

The indirect methods are based on the determination of the CO₂ concentration gradient in the soil (Camarda et al. 2006). These methods can be applied only if the transport of the gas is dominated by the diffusion and some properties of the medium are known. Direct methods require dynamic or static procedures, whether or not a flux of air is used to extract gas from the soil. The dynamic procedures require some corrections depending on the physical properties of

the soil in the measurement point and on the design of the instrumental apparatus. Furthermore, all dynamic procedures are affected by overpressurization or depressurization depending upon the magnitude of the air flux chosen by the operator and according to Kanemasu et al. (1974) results are strongly affected by the physical modifications induced by pumping under different flux regimes. Camarda et al. (2006) present a demonstration of the indirect method applied to measuring CO₂ fluxes from Vulcano (Italy), and show that with low pumping rates the sensitivity of the method to soil permeability is reduced.

The accumulation chamber method (or closed-chamber method) is a direct, static method originally used in agricultural sciences to determine soil respiration (Parkinson 1981) and then successfully adapted to measure CO₂ soil flux of volcanological interest by Tonani and Miele (1991). This method is based on the measurement of the CO₂ concentration increase inside an open-bottomed chamber of known volume, inverted on the soil surface. The initial rate of change of the concentration is proportional to the CO₂ flux (Tonani and Miele 1991; Chiodini et al. 1996). The method does not require either assumptions about soil characteristics or the regime of the flux (advective/diffusive). The method has been tested by several authors under controlled laboratory conditions and provides reproducibility of 10% (Chiodini et al. 1998). In a field reproducibility test of the method, carried out at two points with high and low CO₂ flux, Carapezza and Granieri (2004) found an uncertainty of 12% for high fluxes and 24% for low fluxes. Multiple measurements performed by ground-based methodologies allow a mapping of CO₂ flux and an estimation of the total CO₂ release by use of interpolation algorithms (Cardellini et al. 2003; Chiodini et al. 2005).

Eddy covariance or alternately Eddy correlation (EC) is a micrometeorological technique (e.g., Baldocchi 2003) recently proposed as a method to monitor volcanic CO₂ emissions (Werner et al. 2000, 2003; Anderson and Farrar 2001; Lewicki et al. 2008). The basis of the EC is the calculation of the flux at the surface through the covariance between the fluctuations of the vertical component of the wind and the fluctuations of the gas concentration in atmosphere. The EC provides advantage of being an automated, time-averaged and area-integrated technique with a spatial scale significantly larger (square meter to square kilometer) than that of the ground-based methods (e.g., accumulation chamber). However the volcanic environment is often too heterogeneous for EC application, as suggested by the theory underlying EC, in terms of spatial and temporal variability of surface fluxes and morphology of the measuring field.

Diffusive degassing of deep CO₂ in tectonically active areas

Since the early work of Irwin and Barnes (1980), it has become clear that a close relationship exists between active tectonic areas and anomalous crustal emissions of CO₂. Due to their high crustal permeability, faults act as preferential pathways for the upward migration and eventual release of deep gases to the aquifers or directly to the atmosphere. Regional aquifers located in areas of high CO₂ flux can dissolve most or part of the deeply generated gas because the relatively high solubility of CO₂ in water. A carbon mass balance in the involved aquifer can be used to obtain an estimation of the amount of CO₂ dissolved by groundwater. However, the large range of ¹³C_{CO₂ observed in such aquifers suggests that carbon can derive from multiple sources: atmospheric C, biogenic C, carbonate minerals derived C and deeply derived C. Therefore, an approach by coupling groundwater chemistry with hydrologic and isotopic data has to be applied in order to differentiate shallow versus deep sources. Chiodini et al. (2004) showed that in the tectonically active area of the Italian Apennines, approximately 40% of the inorganic carbon in the groundwater derives from magmatic sources. This observation suggests that there may be significant amounts of magmatic CO₂ released in tectonic areas, perhaps a similar order of magnitude as subaerial volcanic degassing (Chiodini et al. 2004). Indeed, in volcanic areas, the dissolved CO₂ in groundwater can be a significant component of the total CO₂ flux at the volcano (e.g., Rose and Davisson 1996; Sorey et al. 1998; Inguaggiato et al. 2012).}

Submarine measurements

Volatile emissions from the axis of mid-ocean ridges (MORs) in the form of black smokers are dramatic examples of submarine deep carbon emissions, however they tend to be short-lived and unpredictable, making collection of gas samples challenging. Nevertheless measurements of the composition and flux of such emissions have been conducted at dozens of sites (Kelley et al. 2004), using a wide range of gas collection techniques from piloted and remotely controlled submersible craft, allowing later analysis of gas samples in the laboratory. Direct measurements have been performed on only a fraction of the world's MOR, and therefore previous work on the fluxes of CO₂ from MOR has focused on quantifying emissions relative to a better-constrained global production parameter. These have included crustal production rates and the C content of the mantle (Gerlach 1989, 1991; Javoy and Pineau 1991; Holloway 1998; Cartigny et al. 2001; Saal et al. 2002), global mantle ³He flux (Corliss et al. 1979; Des Marais and Moore 1984; Marty and Jambon 1987; Sarda and Graham 1990; Graham and Sarda 1991; Marty and Zimmerman 1999), hydrothermal fluid flux (Elderfield and Schultz 1996) and the CO₂/³He ratio in hydrothermal plumes. The latter two require estimates of global ³He fluxes, which are produced primarily at MORs (Allard et al. 1992).

An important aspect of MOR volcanism is that during the process of formation CO₂ reacts with hot rock, sequestering CO₂. In addition, dissolved carbonate in seawater reacts progressively within the shallowest ~60 m oceanic crust, producing steadily higher carbonate concentrations with increasing crustal age. Measurements of drill cores of the upper oceanic crust allowed Alt and Teagle (1999) to quantify the magnitude of the CO₂ sink produced by crust reactions as of the magnitude of 150 Mt/yr CO₂. This is of similar magnitude to the MOR CO₂ flux of 97 ± 40 Mt/yr CO₂, indeed it is probable that reactions in the oceanic crust absorb more CO₂ than is emitted from MORs.

As well as emissions from the main axis of the MOR, degassing takes place on the flanks of the ridge, driving circulation of seawater through the crust. Sansone et al. (1998) sampled gas emissions from the eastern flank of the Juan de Fuca ridge with the Alvin deep sea vessel, both directly and through inverted funnels to concentrate the gas flow into titanium gas-tight samplers (Massoth et al. 1989). Further samples were collected using titanium syringe samples (Von Damm et al. 1985). Gas samples were acidified and then vacuum-extracted at sea with a glass/stainless-steel vacuum line. The total gas volume was determined with high precision capacitance manometers, and the extracted gas from each sample was sealed in break-seal glass ampoules for analysis ashore with gas chromatography and mass spectrometry.

Gas emissions also occur from active submarine arc volcanoes. Lupton et al. (2008) measured gas output from eleven volcanoes along the Mariana and Tonga-Kermadec arcs with remote controlled vehicles during three expeditions. Four of these volcanoes were found to produce distinct gas and liquid CO₂ gas emissions, together with hydrothermal emissions from the main vents. Vent emissions were sampled using seawater-filled titanium alloy gas-tight bottles connected via tubing to a Ti sampling snout inserted directly into the vent. A valve was then opened and hydraulic pressure filled the bottle with a sample of vent gas. On the ship, samples were acidified and transferred under vacuum ampoules made of Pyrex and low-He permeability aluminosilicate glass for later laboratory analysis. Collection of liquid CO₂ droplets was challenging due to the ~1000 fold expansion of liquid CO₂ when converted to CO₂ gas at 1 atm pressure, necessitating the use of a small volume Ti gas-tight bottle. Gas bubbles were collected with a plastic cylinder with relief valve that was placed over the emission until filled.

Understanding of the carbon balance in a subduction zone requires knowledge of the amount of carbon entering the zone within the subducting slab, CO₂ loss from main arc volcanism and back arc, and the submarine fore-arc (see Fig. 1). This latter was measured

from seeps in the Central America subduction zone by Füri et al. (2010). Seep fluids were collected over a 12 month period at the submarine segment of the Costa Rica fore-arc margin using 1/8 inch diameter copper tubing attached to a submarine flux meter operating in continuous pumping mode to measure CO₂ and CH₄ fluxes. Temporal variations during the sampling period were revealed by cutting the copper tubing in 0.4 m sections under vacuum and extracting the stored volatile samples in the laboratory for isotope ratios and compositions.

Volcanic lakes are significant, but previously unrecognized (Pérez et al. 2011) contributors to global deep CO₂ budgets. CO₂ gas emissions from volcanic lakes are in the form of both diffuse degassing from the lake surface and bubbling (Mazot and Taran 2009). Lake CO₂ emissions were therefore measured with a floating gas accumulation chamber with an in-built NIR sensor to measure CO₂ concentrations. Conversion of CO₂ concentrations to fluxes was made using simultaneous measurements of pressure and temperature.

REPORTED MEASUREMENTS OF DEEP CARBON FLUXES

Subaerial volcanism

During the Holocene, ~1500 volcanoes on land erupted and in recorded history there have been 550 known eruptions. Typically 50-70 volcanoes erupt explosively each year and ~500 produce a gas emission either through hydrothermal systems or open-vent degassing (Siebert and Simkin 2002). Of these, only a small fraction have had their CO₂ flux measured directly, however the number of measured volcanoes has greatly increased in recent years. In this section we first present published data on measured volcanic CO₂ emission rates from active volcanoes, diffuse degassing of volcanic areas and tectonically active areas, followed by volcanic lakes and submarine emissions. We conclude by producing a global sum of CO₂ emission rates.

We report in Table 1 all known volcanic plume CO₂ flux measurements from persistently degassing volcanoes. We have chosen data in which CO₂ fluxes were measured either by near simultaneous measurement of CO₂/SO₂ ratios and SO₂ / gas flux, or direct measurement of the CO₂ flux. This was done because the number of volcanoes for which CO₂ flux has been measured accurately has greatly increased in the last years, with 40 new measurements reported since 2000. We performed a simple average of the reported CO₂ flux measurements for each volcano to produce Table 2, a summary which allows the total CO₂ flux from 33 measured volcanic gas plumes to be calculated as 59.7 Mt/yr. We note that this measured flux by itself is higher than the maximum estimated for global passive degassing from Williams et al. (1992), highlighting the fundamental importance of direct measurements of volcanic CO₂ fluxes in quantifying the volcanic CO₂ inventory.

In Table 3, we report diffuse CO₂ fluxes from historically active volcanoes, which have been verified through isotopic analysis to be of magmatic origin. This list is not an exhaustive collation of all diffuse CO₂ degassing measurements, but reflects the most updated or complete diffuse CO₂ flux measured at each volcano. The total CO₂ flux from the 30 measured volcanoes is 6.4 Mt/yr, including emissions from diffuse soil degassing and those measured in groundwater.

CO₂ fluxes from tectonic structures, hydrothermal systems or inactive volcanic areas are reported in Table 4, distinguishing between measured (or estimated) fluxes from soils from those dissolved in groundwaters. The two major measured contributors to the total tectonic, hydrothermal and inactive volcano CO₂ flux of 66 Mt/yr are tectonic degassing in Italy (10 Mt/yr, Chiodini et al. 2004) and hydrothermal emissions from Yellowstone (8.6 Mt/yr, Werner and Brantley 2003). We also include in this list an estimate of the total CO₂ flux produced by hydrothermal activity in Indonesia-Philippines (1.8 Mt/yr, Seward and Kerrick 1996) and

Table 1. Volcanic plume CO₂ fluxes from persistently degassing volcanoes
(alphabetically ordered by country)

Volcano	Location	CO₂ Flux (t/d)	CO₂ Flux (Mt/yr)	Method	Date	Reference
Erebus	Antarctica	1,930	0.70	Airborne Li-COR	Dec 1997, Dec 1999, Jan 2001	Wardell et al. (2004)
Erebus	Antarctica	1,330	0.49	Ground-based OP-FTIR and scanning UV spectrometer	Dec 2004	Oppenheimer & Kyle (2008)
Villarrica	Chile	477	0.17	Ground-based OP-FTIR and airborne UV spectrometer traverses	Mar 2009	Sawyer et al. (2011)
Galeras	Colombia	1,020	0.37	High temp. summit fumarole sampling and COSPEC	1989-1995	Zapata et al. (1997)
Nyiragongo	DR Congo	95,500	34.86	Direct sampling with video footage of plume dimensions and rise rate	1959, 1972	Le Guern (1987)
Nyiragongo	DR Congo	9,320	3.40	Ground-based OP-FTIR and UV spectrometer vehicle traverses	May/June 2005, Jan 2006	Sawyer et al. (2008a)
Sierra Negra	Ecuador (Galápagos)	394	0.14	Ground-based multi-gas sensor and UV spectrometer walking traverses	June-July 2006	Padron et al. (2012)
Erta Ale	Ethiopia	54	0.02	Based on heat budget calculations	1971, 1973, 1974	Le Guern et al. (1979)
Erta Ale	Ethiopia	60	0.02	Ground-based OP-FTIR and UV spectrometer walking traverses	15 Oct 2005	Sawyer et al. (2008b)
Grimsvötn	Iceland	532	0.19	Sampling of subglacial crater lake	1954-1991	Agustsdottir & Brantley (1994)
Merapi	Indonesia	240	0.09	High temp. dome fumarole sampling and routine SO ₂ flux measurements		Toutain et al. (2009)
Etna	Italy	35,000	12.8	Plume sampling or airborne IR analysers and ground-based and airborne COSPEC;	1977-1984	Allard et al. (1991)

Table 1 (continued). Volcanic plume CO₂ fluxes

Volcano	Location	CO ₂ Flux (t/d)	CO ₂ Flux (Mt/yr)	Method	Date	Reference
Etna	Italy	9,000	3.29	Ground-based multi-gas (Voragine and NE) and UV spectrometer vehicle traverses	Sept 2004- Sept 2005	Aiuppa et al. (2006)
Etna	Italy	5,090	1.86	Ground-based multi-gas (Voragine) and UV spectrometer vehicle traverses	May 2005- Nov 2006	Aiuppa et al. (2008)
Stromboli	Italy	4,350	1.59	Fumarole and quiescent plume sampling and airborne COSPEC	1980-1993	Allard et al. (1994)
Stromboli	Italy	1,073	0.39	Ground-based OP-FTIR automated UV spectrometer network	9 Apr 2002 (IR) 2006 (UV)	Burton et al. (2007a,b)
Stromboli	Italy	550	0.20	Ground-based multi-gas sensor and automated UV spectrometer network	Sept 2008- Jul 2010	Aiuppa et al. (2011)
Vulcano	Italy	420	0.15	Ground-based multi-gas measurement and UV spectrometer vehicle traverses	Dec 2004	Aiuppa et al. (2005)
Vulcano	Italy	170	0.06	UAV mounted electrochemical sensors and UV spectrometer	Apr 2007	McGonigle et al. (2008)
Vulcano	Italy	362	0.13	High temp. fumarole sampling and UV spectrometer vehicle traverses	Sept 2007	Inguaggiato et al. (2012)
Miyakejima	Japan	14,500	5.29	Airborne Li-COR, SO ₂ electrochemical sensor or pulsed fluorescence SO ₂ analyzer and COSPEC	2000-2001	Shinohara et al. (2003)
Satsuma-Iwojima	Japan	100	0.04	High temp. fumarole sampling and COSPEC;	Oct 1999	Shimoike et al. (2002)
Popocatepetl	Mexico	9,000	3.29	Airborne Li-COR ladder surveys	Jun 1995	Gerlach et al. (1997)
Popocatepetl	Mexico	40,000	14.6	Airborne Li-COR ladder surveys	1996-1998	Delgado et al. (1998)
Popocatepetl	Mexico	38,000	13.87	Passive OP-FTIR and COSPEC	Feb 1998	Goff et al. (2001)

Soufrière Hills	Montserrat	1,468	0.54	Ground-based multi-gas sensor and automated UV spectrometer network	Jul 2008	Edmonds et al. (2010)
White Island	New Zealand	950	0.35	High temp. fumarole sampling and airborne COSPEC	1982-1984	Rose et al. (1986)
White Island	New Zealand	2,610	0.95	Airborne Li-COR ladder survey	Jan 1998	Wardell et al. (2001)
Masaya	Nicaragua	2,940	1.07	Ground-based OP-FTIR and COSPEC vehicle traverses	1998-1999	Burton et al. (2000)
Masaya	Nicaragua	930	0.34	Ground-based OP-FTIR and UV spectrometer vehicle traverses	Mar 2009	Martin et al. (2010)
Kudryavy	Russia	50	0.02	Fumarole sampling and ground-based COSPEC	Aug 1995	Fischer et al. (1998)
Bezymianny	Russia	990	0.37	Dome fumarole sampling and FLYSPEC ground-based scanning and airborne traverses	Aug 2007, Jul 2009	Lopez et al. (in press)
Gorely	Russia	660	0.24	Ground-based multi-gas measurement and UV camera	6 Sept 2011	Aiuppa et al. (2012)
Oldoinyo Lengai	Tanzania	6,630	2.42	Airborne Li-COR measurements	Jun 1994	Brantley & Koepnick (1995)
Augustine	USA	1,760	0.64	Restored gas samples and COSPEC	1986-1987	Symonds et al. (1992)
Douglas	USA	trace	trace	Airborne Li-COR measurements	2000-2006	Doukas & McGee (2007)
Griggs	USA	not detected	not detected	Airborne Li-COR measurement	1 Jul 2002	Doukas & McGee (2007)
Iliamna	USA	131	0.05	Airborne Li-COR measurements	1996-2005	Doukas & McGee (2007)
Kīlauea, Pu'u 'Ō'ō	USA	3,950	1.45	Airborne closed-path FTIR	15 Feb 1984, 4 Mar 1984	Greenland et al. (1985)
Kīlauea, Pu'u 'Ō'ō	USA	300	0.11	Airborne Li-COR, closed-path FTIR and COSPEC	19 Sept 1995	Gerlach et al. (1998)
Kīlauea, Pu'u 'Ō'ō	USA	396	0.14	Airborne Visible/Infrared Imaging Spectrometer (AVIRIS)	26 Apr 2000	Spinetti et al. (2008)

Table 1 (continued). Volcanic plume CO₂ fluxes

Volcano	Location	CO ₂ Flux (t/d)	CO ₂ Flux (Mt/yr)	Method	Date	Reference
Kilauea summit	USA	1,600	0.58	Airborne closed-path FTIR	13 Feb 1984	Greenland et al. (1985)
Kilauea summit	USA	8,500	3.10	Ground-based Li-COR, closed-path FTIR and COSPEC	20 Sept 1995, 20 Oct 1998, 6 May 1999	Gerlach et al. (2002)
Kilauea summit	USA	4,900	1.79	Ground-based Li-COR, Interscan electrochemical SO ₂ analyzer and COSPEC;	Jun-Jul 2003	Hager et al. (2008)
Mageik	USA	341	0.12	Airborne Li-COR measurements	2000-2006	Doukas & McGee (2007)
Martin	USA	56	0.02	Airborne Li-COR measurements	1998-2006	Doukas & McGee (2007)
Mt. Baker	USA	187	0.07	Airborne Li-COR measurement	13 Sept 2000	McGee et al. (2001)
Mt. Baker	USA	150	0.05	Airborne Li-COR measurement	2007	Werner et al. (2009)
Peulik	USA	not detected	not detected	Airborne Li-COR measurement	24 May 1998	Doukas & McGee (2007)
Reboubt	USA	18	0.01	Airborne Li-COR measurements	1997-2005	Doukas & McGee (2007)
Spurr	USA	633	0.23	Airborne Li-COR measurements	2004-2006	Doukas & McGee (2007)
Spurr Crater Peak	USA	334	0.12	Airborne Li-COR measurements	1996-2006	Doukas & McGee (2007)
Ukinrek Maars	USA	187	0.07	Airborne Li-COR measurement	24 May 1998	Doukas & McGee (2007)
Veniaminof	USA	not detected	not detected	Airborne Li-COR measurement	2 Aug 2003	Doukas & McGee (2007)
Ambrym	Vanuatu	20,000	7.30	Multi-gas sensor & airborne UV spectrometer traverses	2007	Allard et al. (2009)
Yasur	Vanuatu	840	0.31	Ground-based multi-gas sensor and UV spectrometer vehicle traverses	21 Oct 2007	Métrich et al. (2011)

Table 2. Mean volcanic plume CO₂ fluxes from persistently degassing volcanoes (ordered by CO₂ flux)

Volcano	Country	CO₂ Flux (t/d)	CO₂ Flux (Mt/yr)
Nyiragongo	DR Congo	52,410	19.13
Popocatepetl	Mexico	29,000	10.59
Ambrym	Vanuatu	20,000	7.30
Etna	Italy	16,363	5.97
Miyakejima	Japan	14,500	5.29
Oldoinyo Lengai	Tanzania	6,630	2.42
Kilauea	USA	6,549	2.39
Stromboli	Italy	1,991	0.73
Masaya	Nicaragua	1,935	0.71
White Island	New Zealand	1,780	0.65
Augustine	USA	1,760	0.64
Erebus	Antarctica	1,630	0.59
Soufrière Hills	Montserrat	1,468	0.54
Galeras	Colombia	1,020	0.37
Bezymianny	Russia	990	0.36
Spurr	USA	967	0.35
Yasur	Vanuatu	840	0.31
Gorely	Russia	660	0.24
Grímsvötn	Iceland	532	0.19
Villarrica	Chile	477	0.17
Sierra Negra	Ecuador (Galápagos)	394	0.14
Mageik	USA	341	0.12
Vulcano	Italy	317	0.12
Merapi	Indonesia	240	0.09
Ukinrek Maars	USA	187	0.07
Mt. Baker	USA	169	0.06
Iliamna	USA	131	0.05
Satsuma-Iwojima	Japan	100	0.04
Erta Ale	Ethiopia	57	0.02
Martin	USA	56	0.02
Kudryavy	Russia	50	0.02
Redoubt	USA	18	0.01
Douglas	USA	trace	trace
	Total	163,562	59.70

Table 3. Diffuse CO₂ emissions from historically active volcanoes (alphabetically ordered by country)

Volcano	Country	CO₂ soil gas flux (t/yr)	Dissolved CO₂ flux (t/yr)	Reference
Erebus	Antarctica	14,600		Wardell et al. (2003)
Sierra Negra	Ecuador (Galápagos)	220,825		Padron et al. (2012)
Santa Ana	El Salvador	59,130		Salazar et al. (2004)
Nea Kameni	Greece	5,621		Chiodini et al. (1998)
Nisyros	Greece	24,784		Caliro et al. (2005)
Hengill volcanic system	Iceland	165,345		Hernández et al. (2012)
Krafla geothermal system	Iceland	84,000		Ármannsson et al. (2007)
Reykjanes volcanic sys.	Iceland	12,660		Óskarsson & Fridriksson (2011)
Merapi	Indonesia	78,475		Toutain et al. (2009)
Etna	Italy	1,000,000	250,000	D'Alessandro et al. (1997)
Ischia	Italy	468,940	9,461	Pecoraiano et al. (2005)
Pantelleria	Italy	361,000	34,000	Favara et al. (2001)
Solfatara, Campi Flegrei	Italy	556,260		Chiodini et al. (2001)
Stromboli	Italy	82,125		Carapezza & Federico (2000)
Vesuvio	Italy	55,115		Fronchini et al. (2004)
Vulcano	Italy	41,975	2,190	Inguaggiato et al. (2012)
Miyakejima (Oyama)	Japan	43,618		Hernández et al.(2001a)
Satsuma-Iwojima	Japan	7,300		Shimoike et al. (2002)
Showa-Shinzan	Japan	3,760		Hernández et al. (2006)
Usu	Japan	60,712		Hernández et al.(2001b)
Popocatepetl	Mexico	not detected		Varley & Armienta (2001)
Cerro Negro	Nicaragua	1,022,000		Salazar et al. (2001)
Masaya caldera	Nicaragua	630,720		Pérez et al (2000)
Masaya, Comalito	Nicaragua	6,935		Chiodini et al. (2005)
Rabaul	Papua New Guineau	876,000		Pérez et al. (1998)
Furnas	Portugal		9,358	Cruz et al. (1999)
Oldoinyo Lengai	Tanzania	36,432		Koepenick et al. (1996)
Teide	Tenerife	38,836	64,605	Hernández et al. (2000); Marrero et al. (2008)
Lassen	USA	35,000	7,600	Rose & Davisson (1996)
Mt. Shasta	USA		8,500	Rose & Davisson (1996)
Ukinrek Maars	USA	11,863	1,095	Evans et al. (2009)
	Total (t/yr)	6,004,031	386,809	
	Total (Mt/yr)	6.00	0.39	

Table 4. CO₂ emissions from tectonic, hydrothermal or inactive volcanic areas (alphabetically ordered by country)

Area or vent	Country	CO ₂ soil gas flux (t/yr)	Dissolved CO ₂ flux (t/yr)	Reference
Tengchong Cenozoic volcanic field	China		3,580	Cheng et al. (2012)
Albani Hills	Italy	26,840	157,960	Chiodini & Frondini (2001)
Bossoleto, Siena	Italy	3,500		Mörner & Etiope (2002)
Caldara di Manziana	Italy	73,000		Chiodini et al. (1999)
Campanian degassing structure	Italy		3,080,000	Chiodini et al. (2004)
Castiglioni, Siena	Italy	4,400		Mörner & Etiope (2002)
Latera	Italy	127,750		Chiodini et al. (2007)
Mefite d'Ansanto	Italy	730,000		Chiodini et al. (2010)
Naftia Lake area	Italy	72,217		Giammanco et al. (2007)
Pienza	Italy	4,015		Rogie et al. (2000)
Poggio dell'Ulivo	Italy	73,000		Chiodini et al. (1999)
Rapalano Cecilia	Italy	17,520		Rogie et al. (2000)
Rapalano Mofete Diambra	Italy	35,040		Rogie et al. (2000)
San Sisto	Italy	21,600		Italiano et al. (2000)
Selvena	Italy	6,205		Rogie et al. (2000)
Telese	Italy	20,000		Italiano et al. (2000)
Tuscan Roman degassing structure	Italy		6,160,000	Chiodini et al. (2004)
Umbertide	Italy	5,840		Rogie et al. (2000)
Ustica	Italy	260,000		Etiope et al. (1999)
Hakkoda	Japan	27,010		Hernández et al. (2003)
Taupo	New Zealand		440,000	Seward & Kerrick (1996)
Yangbajain	Tibet	50,370		Chiodini et al. (1998)
Mammoth Mountain	USA	189,800	14,600	Sorey et al. (1998)
Mt Washington and Belknap Crater	USA		2,400	James et al. (1999)
Mt Jefferson	USA		8,000	James et al. (1999)
Mt Bachelor	USA		1,800	James et al. (1999)
Salton Trough	USA		44,000	Kerrick et al. (1995)
Three Sisters	USA		4,400	James et al. (1999)
Yellowstone	USA	8,580,000		Werner & Brantley (2003)
Indonesia-Philippines		1,800,000		Seward & Kerrick (1996)
Subaerial Pacific rim		44,000,000		Seward & Kerrick (1996)
	Total (t/yr)	56,128,107	9,916,740	
	Total (Mt/yr)	56.13	9.92	

the subaerial Pacific rim (44 Mt/yr, Seward and Kerrick 1996). These estimates are based on extrapolations from the CO₂ emissions observed from the 150 km long Taupo Volcanic Zone (New Zealand) to the 18,000 km long Pacific Rim.

The global emissions of CO₂ from volcanic lakes were recently assessed by Pérez et al. (2011), who pointed out that volcanic lakes had not been included in previous estimates of global geological carbon efflux (e.g., Kerrick et al. 2001; Mörner and Etiope 2002). They found that CO₂ emissions increased with increasing acidity in volcanic lakes, reflecting the acidity of the volcanic gas discharge. Measurements were conducted on 32 volcanic lakes which were divided into three types of water based on pH, alkali, neutral and acid. Average flux per unit area for each type was then used to calculate a global volcanic lake estimate, extrapolating to an estimated number of volcanic lakes in the world (769). This number of lakes is greater than the number of lakes reported in the literature (138) by a factor which reflects the regional under-sampling between actual lakes and lakes reported in the scientific literature. This methodology assumes that the average acidity-emission rate relationship in the 32 measured lakes is a faithful average representation of the global lake population. A further estimate was produced by defining 4 populations in the measured data set based on frequency, emission rate and lake size and extrapolating to all 769 volcanic lakes. The combination of these two approaches yielded a global volcanic lake CO₂ emission of 117 ± 19 Mt/yr, of which 94 Mt/yr is attributed to magmatic degassing.

Submarine volcanism

There are three main submarine sources of CO₂, MOR, arc volcanoes and fore-arc degassing (which appears to be dominated by CH₄ emissions (Füri et al. 2010)). Global emissions from MOR have been determined by various authors, as reported in Table 5. The large spread of MOR fluxes, from 4.4 to 792, reflects uncertainties in the dissolved contents of C and global ³He fluxes. Marty and Tolstikhin (1998) performed a careful examination of the CO₂/³He ratios used and determined a median value of 2.2×10^9 with standard deviation of 0.7×10^9 . Using a ³He flux of 1000 ± 250 mol/yr (Farley et al. 1995) they derived a MOR CO₂ flux of 97 ± 40 Mt/yr CO₂. The more recent determination of MOR CO₂ flux from Resing et al. (2004) who measured CO₂/³He in MOR hydrothermal plumes of 55 ± 33 Mt/yr is in reasonable agreement with that estimate.

While measurements of CO₂ release from the cooler flanks of MORs and submarine arc volcanoes increase in number each year, global estimates of submarine CO₂ emissions are extremely difficult to make. The large areal extent and our relatively poor knowledge of the submarine surface suggests that there is ample opportunity for unknown or unrecognized active volcanism (e.g., cold liquid CO₂ emissions, Lupton et al. 2008), but at the current time it is not possible to make quantitative estimates of the global CO₂ emissions from such sources.

INVENTORIES OF GLOBAL VOLCANIC DEEP CARBON FLUX: IMPLICATIONS FOR THE GEOLOGICAL CARBON CYCLE

Estimates of global deep carbon emission rates

In Table 6, we summarize the measured fluxes from the subaerial sources and MOR, and attempt to extrapolate from these measurements to global estimates of the CO₂ flux for each source. In the case of volcanic plume passive degassing the GVN catalogue (Siebert and Simkin 2002) indicates that there are ~150 such actively degassing volcanoes on Earth. While our catalogue of 33 CO₂ flux plume measurements (Table 2: total flux 59.7 Mt/yr) is significantly larger than previously collated, it reflects only 22% of the total number of active volcanoes. While our current compilation includes some large emitters, suggesting that the major sources have been already identified, we highlight how this total flux has increased due

Table 5. MOR global CO₂ flux (Mt/yr)

Method	Min	Max	Reference
Crustal production rates and C content of mantle	10	35	Gerlach (1989)
	22	39.6	Gerlach (1991)
	572	748	Javoy & Pineau (1991)
	128	255	Holloway (1998)
	176	792	Cartigny et al. (2001)
	28.6	41	Saal et al. (2002)
CO ₂ / ³ He in MORB glass, global mantle ³ He flux	75	119	Marty & Jambon (1987)
	18	44	Sarda & Graham (1990)
	106	264	Graham & Sarda (1991)
	66	119	Marty & Zimmerman (1999)
CO ₂ / ³ He in MOR fluids, global mantle ³ He flux	44	70	Corliss et al. (1979)
	20	57	Des Marais & Moore (1984)
Hydrothermal fluid flux and composition	4.4	53	Elderfield & Schultz (1996)
CO ₂ / ³ He in plumes	22	88	Resing et al. (2004)
Summary	4.4	792	

Table 6. Summary of measured volcanic CO₂ fluxes and estimated global emissions (Mt/yr)

Source	Measured CO ₂ flux	N ^o measured	N ^o global	% global	Estimated Global CO ₂ flux	Ref.
Volcanic plume passive degassing	59.7	33	~150	22	271	[1]
Diffuse emissions from historically active volcanoes	6.4	30	~550	5.5	117	[1]
Emissions from tectonic, hydrothermal or inactive volcanic areas	66	—	—	—	>66	[1]
Volcanic lakes	6.7	32	769	4.2	94	[2]
MOR	97	—	—	—	97	[3]
			Total		637	
			Total (no MOR)		540	

References: [1] This work; [2] Perez et al. (2011); [3] Marty and Tolstikhin (1998)

to previously unrecognized large emissions from e.g., Ambrym (Vanuatu). Further large, but as yet unquantified, sources of CO₂ emissions may be present in Papua New Guinea, the Banda Sunda arc and the Vanuatu island chain. We therefore conclude that the clearest and probably most accurate way to extrapolate from the current catalogue of plume emissions to a global estimate is through a linear extrapolation. Extrapolating from the measured 33 to an estimated 150 plume-creating, passively degassing volcanoes we estimate that the global plume CO₂ flux is ~271 Mt/yr (see Table 6).

The total number of historically active volcanoes reported by GVN is ~550, and 30 (5.5%) of these have had diffuse CO₂ soil degassing fluxes quantified, as reported in Table 3, for a total of 6.4 Mt/yr. Extrapolating to a global flux, assuming a similar distribution of fluxes in the unmeasured fluxes as seen in those measured, produces a total of 117 Mt/yr from diffuse degassing from the flanks of historically active volcanoes (Table 6).

The total diffuse CO₂ flux from inactive volcanoes, hydrothermal and tectonic structures reported in Table 4 is more challenging to extrapolate to a global scale. Our current constraints on the tectonic CO₂ flux comes almost entirely from the work of Chiodini et al. (2004) who examined actively degassing tectonic structures in Italy. The abundance of such structures on Earth is unknown, and this therefore represents a source of great uncertainty in estimates of total deep carbon flux. This uncertainty makes it challenging to sensibly extrapolate to a global estimate of tectonic CO₂ fluxes, and therefore we use only reported fluxes, and highlight the possibility that the true total may be significantly larger. Emissions from hydrothermal systems estimated by Seward and Kerrick (1996) are already extrapolated to cover a significant proportion of the volcanically active surface of the Earth. We therefore use the total presented in Table 4 for the CO₂ emissions from tectonic, hydrothermal and inactive volcanoes as a lower limit for the global emission of CO₂ from these sources.

Summing the extrapolated passive plume, diffuse degassing, lake degassing global estimates and emissions from inactive, hydrothermal and tectonic structures produces a total subaerial volcanic flux of 540 Mt/yr, and a global emission (including MOR emissions) of 637 Mt/yr (Table 6). Thus global volcanic CO₂ fluxes are only ~1.8% of the anthropogenic CO₂ emission of 35,000 Mt per year (Friedlingstein et al. 2010).

Comparison with previous estimates of subaerial volcanic CO₂ flux

There have been several papers which estimate the global CO₂ flux, as shown in Table 7. Our update of the global volcanic CO₂ flux, 637 Mt/yr, is larger than the maximum suggested by Marty and Tolstikhin (1998) of 440 Mt/yr. This is in part because CO₂ emissions from volcanic lakes were not addressed in that work. The total subaerial flux we calculate of 540 Mt/year is also higher than that proposed by Mörner and Etiope (2002), due primarily to the improvement in measurements of persistently degassing volcanoes. We note that Mörner and Etiope (2002) included the fluxes from single eruptive events from Pinatubo (1991) and Mt. St. Helens (1980) in their inventory of volcanoes contributing to the annual global CO₂ flux. Other papers cited in Table 7 appear to have significantly underestimated the global subaerial CO₂ flux, primarily due to a lack of field measurements.

Table 7. Global volcanic subaerial CO₂ flux (Mt/yr)

CO ₂ flux	Reference
79	Gerlach (1991)
145	Varekamp et al. (1992)
66	Allard (1992)
88	Marty and Le Cloarec (1992)
65	Williams et al. (1992)
136	Sano and Williams (1996)
242	Marty and Tolstikhin (1998)
99	Kerrick (2001)
300	Mörner and Etiope (2002)
540	This work

Balancing CO₂ emission rates with weathering and subduction rates

In the absence of a continual supply of CO₂ from volcanic and tectonic degassing the CO₂ content of the atmospheres and oceans would be gradually depleted through CO₂ removal by weathering (Gerlach 1991). The fact that, instead, pre-industrial CO₂ concentrations are relatively stable suggest a balance between CO₂ removal by weathering and CO₂ supply by Earth degassing on timescales of ~0.5 Ma (Walker et al. 1987; Berner 1991). Over such timescales weathering of carbonates has no impact on removal of atmospheric CO₂, because

HCO_3^- supplied to the ocean by carbonate weathering releases the CO_2 it captured from the atmosphere during calcite precipitation (Berner 1991). Therefore in the long timescale of the geological carbon cycle CO_2 emissions from geological sources should balance consumption from silicate weathering and oceanic crust alteration. Gaillardet et al. (1999) found that CO_2 consumption from continental silicate weathering was 515 Mt/yr, which matches well with our estimates of subaerial volcanic CO_2 degassing (540 Mt/yr). However, inclusion of 300 Mt/yr CO_2 released by metamorphism (Morner and Etiope 2002) produces a total lithospheric subaerial CO_2 emission of 840 Mt/yr. This is larger than the current estimates of silicate weathering, suggesting that, assuming steady-state, weathering rates might be slightly higher to absorb all the emitted CO_2 .

Dasgupta and Hirschmann (2010) calculated the total ingassing of CO_2 into subduction zones from the combination of three lithologies in the subducting slab, altered oceanic crust, sediments and mantle, producing an estimated CO_2 consumption rate of 403 Mt/yr. We note that this is lower than the CO_2 consumption rate due to silicate weathering (515 Mt/yr, Gaillardet et al. 1999), which is slightly inconsistent (but within uncertainties of such estimates), as the eventual destiny of CO_2 consumed by silicate weathering will be deposition on the seafloor or precipitation within the oceanic crust. In order to maintain steady-state quantities of CO_2 in the exosphere this consumption should be balanced by the total emission from MORs, subaerial degassing and metamorphism (calculated here to be 937 Mt/yr). Given the likely underestimate in our total lithospheric CO_2 emissions arising from lack of knowledge of tectonic degassing, it appears reasonable to conclude that the ingassing rate may be an underestimate. However, it is clear that there are large uncertainties in both sums.

THE ROLE OF DEEP CARBON IN VOLCANIC ACTIVITY

Original CO_2 contents of magma

The volatile content of magmas, together with their evolving viscosity and degassing behavior during ascent, helps to determine whether a volcano will be quietly degassing or violently erupting for a given magma input rate. Models of magma dynamics require knowledge of original volatile content in order to reproduce physically accurate processes occurring during an eruption and in quiescent phases. Furthermore, knowledge of the original volatile contents of magmas allows calculation of the magma mass required to produce an observed gas flux, permitting quantitative comparison of fluxes with geophysical and volcanological observations. Measurements of original volatile contents are therefore of great interest.

Melt inclusions (MIs) provide records of original volatile contents, through analysis of pockets of melt trapped inside growing crystals during magma ascent or storage, and such studies have been carried out on many eruption products. However, the presence of a separate fluid phase at the moment of inclusion entrapment will produce an underestimate in the concentrations of dissolved volatiles. Wallace (2005) concluded that no melt inclusions sample arc magmas undegassed with respect to CO_2 . Blundy et al. (2010) used MI measurements from Mt. St. Helens to show that the dissolved volatile contents of shallow magmas were strongly affected by CO_2 -rich fluids rising from magmas at greater depth, concluding that the original CO_2 contents of arc magmas was likely to be significantly higher than that recorded in MIs. Using inferred CO_2 contents in arc andesites and dacites of 1.5 wt% they calculate that the CO_2 contents of parental, mantle-derived basalts would contain 0.3 wt%. This relatively high CO_2 content is in agreement with previous estimates of volatile contents of arc magmas (Wallace 2005).

Original CO_2 contents of magmas can also be estimated by assuming a steady-state condition for a persistently degassing volcano, and comparing the observed CO_2 flux together with the flux of a more soluble gas species whose original volatile content has been well-characterized, such as SO_2 . Gerlach et al. (2002) performed such a calculation for Kilauea

volcano, Hawai'i, concluding that the bulk CO₂ concentration required to match magma input rates and CO₂ output rates was 0.70 wt%. Dixon and Clague (2001) measured a dissolved CO₂ content of at a depth of 1500-2000 m at the Loihi seamount in the Hawai'i chain. While the CO₂ concentrations were very low, the samples contained fluid-filled vesicles with high CO₂ contents, which allowed a bulk CO₂ concentration of up to 0.63 wt% to be determined, in fair agreement with the estimate from Gerlach et al. (2002). A recent study (Barsanti et al. 2009) introduced a more complex note to the examination of original CO₂ contents at Hawai'i, with a statistical analysis of MI CO₂ and H₂O concentrations which revealed distinct magma batches some of which could contain 2-6 wt% CO₂. High CO₂ contents of magmas feeding Etna and Stromboli have also been proposed based on degassing mass balance calculations, with amounts ranging between 1.6 and 2.2 wt% (Spilliaert et al. 2006; Burton et al. 2007a). These few available estimates of original CO₂ contents from mass balance determinations open the possibility that CO₂ contents of magmas feeding active volcanoes are in general higher than is expected based on CO₂ contents of melt inclusions.

Importance of a deep exsolved volatile phase on magma dynamics and eruptive style

The presence of a CO₂-rich volatile phase at great pressure can strongly affect the dynamics of magma ascent and eruption, because the style and intensity of eruptive activity is controlled in part by the distribution of gas phase in a magma during eruption (Eichelberger et al. 1986; Jaupart and Vergnolle 1988). Persistently degassing volcanoes can release vast amounts of gas at the surface non-explosively, implying storage within the crust of large volumes of degassed magma (Crisp 1984, Francis et al. 1993) via magma convection (Kazahaya et al. 1994).

Exsolved volatiles can ascend from depth, accumulating in foams that can produce Strombolian activity (e.g., Menand and Phillips 2007; Jaupart and Vergnolle 1988). Gas can stream through magma from depth to the surface (Wallace et al. 2005), as surmised to occur at Soufrière Hills volcano, Montserrat (Edmonds et al. 2010) and Stromboli volcano (Aiuppa et al. 2010). Perhaps most importantly of all however, exsolved gas accumulation can produce powerful explosive eruptions. The eruption of Pinatubo in 1991 (Pallister et al. 1992) was one of the most violent in recorded history. It produced a much greater mass of S than was to be expected from dissolved S contents and the volume of erupted material, suggesting the presence of a voluminous pre-eruptive gas phase (Wallace and Gerlach 1994), likely produced from basaltic underplating crystallizing as anhydrite (Matthews et al. 1992) which triggered the eruption (Pallister et al. 1992).

MAGNITUDE OF ERUPTIVE DEEP CARBON EMISSIONS

It is useful to compare the CO₂ emission rates for subaerial volcanism of 540 Mt/yr reported in Table 7 with a single large eruption such as the ~5 km³ eruption of Mt. Pinatubo in 1991, producing ~50 Mt of CO₂ (Gerlach et al. 2011), equivalent to merely ~5 weeks of global subaerial volcanic emissions. The Pinatubo 1991 syn-eruptive emission is therefore dwarfed by the time-averaged continuous CO₂ emissions from global volcanism. Indeed, the present day CO₂ emission rate from the lake filling the crater formed during the eruption of Pinatubo is 884 t/d (Perez et al. 2011), suggesting that in the 31 years since that eruption ~10 Mt of CO₂ has been produced, ~20% of that emitted during the eruption.

Using the volumes of erupted material produced by the three largest eruptions of the last 200 years (Self et al. 2006) we may estimate their CO₂ emissions, assuming a similar erupted volume to CO₂ emission amount to that estimated for Pinatubo (10 Mt CO₂ per km³ erupted, equivalent to ~1 wt% CO₂ content). The eruption of Tambora (Indonesia) in 1815 is estimated to have produced 30 km³ of products (Self et al. 2006), with an inferred output of 300 Mt of CO₂. Krakatua in 1883 (Indonesia) and Katmai-Novarupta in 1912 (Alaska) each produced 12

km³, and ~120 Mt of CO₂. The total CO₂ output of the four largest eruptions in the last 200 years is therefore ~600 Mt of CO₂, slightly less than we estimate for subaerial volcanic degassing in a single year, and therefore only 0.6% of the amount of gas released through continuous volcanic activity in the same time period. It therefore appears that the continuous degassing of active and inactive volcanoes dominates the short-lived paroxysmal emissions produced in large eruptions.

Crisp (1984) calculated that the average eruption rate from volcanoes over the last 300 years was 0.1 km³ magma per year, which with ~1 wt% CO₂ content suggests an annual output of ~1 Mt CO₂, only 0.2% of the estimated annual subaerial CO₂ emissions. This demonstrates that degassing of unerupted magma dominates degassing of erupted lava on the planet, and emphasizes the fundamental role that unerupted magmatic intrusions must have in contributing to the global volcanic CO₂ flux. Such intrusions may produce unexpectedly high CO₂ emissions if they interact with crustal carbonates (Troll et al. 2012). This important process could be assessed quantitatively if a method could be developed for measuring volcanic ¹²C/¹³C ratios in the field.

SUMMARY

In recent years, measurements of CO₂ flux from volcanoes and volcanic areas have greatly increased, particularly on persistently degassing volcanoes, of which ~22% have had their CO₂ flux quantified. Notwithstanding this progress, it is clear that the CO₂ emissions from the majority of volcanic sources are still unknown. Using the available data from plume measurements from 33 degassing volcanoes we determine a total CO₂ flux of 59.7 Mt/yr. Extrapolating this to ~150 active volcanoes produces a total of 271 Mt/yr CO₂. Extrapolation of the measured 6.4 Mt/yr of CO₂ emitted from the flanks of 30 historically active volcanoes to all 550 historically active volcanoes produces a global emission rate of 117 Mt/yr. Perez et al. (2011) calculated the global emission from volcanic lakes to be 94 Mt/yr CO₂. The sum of these fluxes produces an updated estimate of the global subaerial volcanic CO₂ flux of 474 Mt/yr. Emissions from tectonic, hydrothermal and inactive volcanic areas contribute a further 66 Mt/yr to this total (Table 6), producing a total subaerial volcanic emission of 540 Mt/yr. An extrapolation to a global estimate is not straightforward for tectonic-related degassing, as the number of areas which produce such emissions is not known. Given the fact that ~10 Mt/yr is produced by Italy alone it is possible that the global total is significant, and this merits further investigation. We highlight also that the magnitude of CO₂ emissions from both cold and hot non-MOR submarine volcanic sources are currently effectively unknown.

Our subaerial volcanic CO₂ flux matches well with estimates of CO₂ removal rates of 515 Mt/yr due to silicate weathering, which, over timescales of 0.5 Ma, should balance lithospheric CO₂ emissions. However, inclusion of the metamorphic CO₂ flux of 300 Mt/yr calculated by Morner and Etiope (2002) produces a total subaerial lithospheric flux of 840 Mt/yr, suggesting that, assuming steady-state, weathering rates might be slightly higher in order to absorb all the CO₂ emitted from the lithosphere.

The global subaerial CO₂ flux we report is higher than previous estimates, but remains insignificant relative to anthropogenic emissions, which are two orders of magnitude greater at 35,000 Mt/yr (Friedlingstein et al. 2010). Nevertheless, it is clear that uncertainties in volcanic CO₂ emission rates remain high and significant upward revisions of the lithospheric CO₂ flux cannot be ruled out. This uncertainty also limits our understanding of global volcanic carbon budgets and the evolution of the distribution of CO₂ between the crust and the mantle. Furthermore, with the notable exception of continuous CO₂ flux monitoring at a handful of volcanoes we have very little data with which to assess CO₂ flux variations across different timescales. It is clear that there is much further work to be done surveying CO₂ emissions from both active and inactive volcanoes.

Continuous global CO₂ emissions from passively degassing volcanoes over timescales longer than a few months dominate CO₂ emissions produced by relatively short-lived eruptions. Nevertheless, we highlight that dramatic CO₂ emissions may occur during magmatic intrusion events, and that the sporadic and short-term nature of field measurements to date may lead to such events being missed. To this end, robust field-portable instruments capable of measuring ¹²C/¹³C ratios in volcanic CO₂ emissions would be of great utility in order to distinguish CO₂ produced during metamorphism of crustal carbonates from magmatic CO₂.

ACKNOWLEDGMENTS

We thank Adrian Jones and Bob Hazen for the opportunity to contribute to this volume. Adrian Jones' and Tamsin Mather's constructive reviews of an earlier version of this work are greatly appreciated. Sara Barsotti is thanked for her simulations of CO₂ concentrations in the Etna plume with VOLCALPUFF. MB acknowledges support from ERC project CO2Vole 279802.

REFERENCES

- Ágústsdóttir AM, Brantley SL (1994) Volatile fluxes integrated over four decades at Grimsvötn volcano, Iceland. *J Geophys Res* 99:9505-9522, doi: 10.1029/93JB03597
- Aiuppa A, Bertagnini A, Metrich N, Moretti R, Di Muro A, Liuzzo M, Tamburello G (2010) A model of degassing for Stromboli volcano. *Earth Planet Sci Lett* 295(1-2):195-204
- Aiuppa A, Burton M, Allard P, Caltabiano T, Giudice G, Gurrieri S, Liuzzo M, Salerno G (2011) First observational evidence for the CO₂-driven origin of Stromboli's major explosions. *Solid Earth* 2(2):135-142, doi: 10.5194/se-2-135-2011
- Aiuppa A, Federico C, Giudice G, Gurrieri S (2005) Chemical mapping of a fumarole field: La Fossa Crater, Vulcano Island (Aeolian Islands, Italy). *Geophys Res Lett* 32:L13309, doi: 10.1029/2005GL023207
- Aiuppa A, Federico C, Giudice G, Gurrieri S, Liuzzo M, Shinohara H, Favara R, Valenza M (2006) Rates of carbon dioxide plume degassing from Mount Etna volcano. *J Geophys Res Solid Earth* 111:B09207, doi: 10.1029/2006JB004307
- Aiuppa A, Giudice G, Gurrieri S, Liuzzo M, Burton M, Caltabiano T, McGonigle AJS, Salerno G, Shinohara H, Valenza M (2008) Total volatile flux from Mount Etna. *Geophys Res Lett* 35(24):L24302, doi: 10.1029/2008GL035871
- Aiuppa A, Giudice G, Liuzzo M, Tamburello G, Allard P, Calabrese S, Chaplygin I, McGonigle AJS, Taran Y (2012) First volatile inventory for Gorely volcano, Kamchatka. *Geophys Res Lett* 39:L06307, doi: 10.1029/2012GL051177
- Aiuppa A, Moretti R, Federico C, Giudice G, Gurrieri S, Liuzzo M, Papale P, Shinohara H, Valenza M (2007) Forecasting Etna eruptions by real-time observation of volcanic gas composition. *Geology* 35:1115-1118, doi: 10.1130/G24149A.1
- Allard P (1992) Global emissions of helium-3 by subaerial volcanism. *Geophys Res Lett* 19:1479-1481
- Allard P, Aiuppa A, Bani P, Metrich N, Bertagnini A, Gauthier PG, Parello F, Sawyer GM, Shinohara H, Bagnato E, Mariet C, Garaebiti E, Pelletier B (2009) Ambrym basaltic volcano (Vanuatu Arc): volatile fluxes, magma degassing rate and chamber depth. *AGU Fall Meeting 2009*, abstract #V24C-04
- Allard P, Burton MR, Mure F (2005) Spectroscopic evidence for a lava fountain driven by previously accumulated magmatic gas. *Nature* 433(7024):407-410, doi: 10.1038/nature03246
- Allard P, Carbone J, Dajlevic D, Le Bronec J, Morel P, Robe MC, Maurenas JM, Faivre-Pierret R, Martin D, Sabroux JC, Zettwoog P (1991) Eruptive and diffuse emissions of CO₂ from Mount Etna. *Nature* 351:387-391, doi: 10.1038/351387a0
- Allard P, Carbone J, Metrich N, Loyer H, Zettwoog P (1994) Sulphur output and magma degassing budget of Stromboli volcano. *Nature* 368(6469):326-330, doi: 10.1038/368326a0
- Alt JC, Teagle DAH (1999) The uptake of carbon during alteration of ocean crust. *Geochim Cosmochim Acta* 63:1527-1535
- Anderson DE, Farrar CD (2001) Eddy covariance measurement of CO₂ flux to the atmosphere from an area of high volcanogenic emissions, Mammoth Mountain, California. *Chem Geol* 177(1-2):31-42, doi: 10.1016/S0009-2541(00)00380-6
- Andres RJ, Kasgnoc AD (1998) A time-averaged inventory of subaerial volcanic sulfur emissions. *J Geophys Res* 103:25251-25261, doi: 10.1029/98JD02091

- Ármansson H, Fridriksson T, Wiese F, Hernández P, Pérez N (2007) CO₂ budget of the Krafla geothermal system, NE-Iceland. *In: Bullen TD, Wang Y (eds) Water-Rock Interaction*. Taylor & Francis Group, London, p 189-192
- Bahcall JN, Pinosmeault MH, Basu S (2001) Solar models: Current epoch and time dependences, neutrinos, and helioseismological properties. *Astrophys J* 555:990, doi: 10.1086/321493
- Baldocchi DD (2003) Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future. *Global Change Bio* 9(4):479-492, doi: 10.1046/j.1365-2486.2003.00629.x
- Barsanti M, Papale P, Barbato D, Moretti R, Boschi E, Hauri E, Longo A (2009) Heterogeneous large total CO₂ abundance in the shallow magmatic system of Kilauea volcano, Hawaii. *J Geophys Res* 114:B12201, doi: 10.1029/2008JB006187
- Barsotti S, Neri A, Scire JS (2008) The VOL-CALPUFF model for atmospheric ash dispersal: 1. Approach and physical formulation. *J Geophys Res* 113:B03208, doi: 10.1029/2006JB004623
- Baubron JC, Allard P, Toutain JP (1990) Diffuse volcanic emissions of carbon dioxide from Vulcano Island, Italy. *Nature* 344(6261):51-53, doi: 10.1038/344051a0
- Berner RA (1991) A model for atmospheric CO₂ over Phanerozoic time. *Am J Sci* 291:339-376, doi: 10.2475/ajs.291.4.339
- Berner RA, Lasaga AC (1989) Modeling the geochemical carbon cycle. *Sci Am* 260:74-81
- Blundy J, Cashman KV, Rust A, Witham F (2010) A case for CO₂-rich arc magmas. *Earth Planet Sci Lett* 290:289-301, doi: 10.1016/j.epsl.2009.12.013
- Brantley SL, Koepnick KW (1995) Measured carbon dioxide emissions from Oldoinyo Lengai and the skewed distribution of passive volcanic fluxes. *Geology* 23(10):933-936, doi: 10.1130/0091-7613(1995)023<0933:MCDEFO>2.3.CO;2
- Brühl C, Lelieveld J, Crutzen PJ, Tost H (2012) The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate. *Atmos Chem Phys* 12:1239-1253, doi: 10.5194/acp-12-1239-2012
- Burton M, Allard P, Murè F, La Spina A (2007a) Depth of slug-driven strombolian explosive activity. *Science* 317:227-230
- Burton MR, Caltabiano T, Mure F, Salerno G, Randazzo D (2009) SO₂ flux from Stromboli during the 2007 eruption: Results from the FLAME network and traverse measurements. *J Volcanol Geotherm Res* 182:214-220, doi: 10.1016/j.jvolgeores.2008.11.025
- Burton MR, Mader HM, Polacci M (2007b) The role of gas percolation in quiescent degassing of persistently active basaltic volcanoes. *Earth Planet Sci Lett* 264:46-60, doi: 10.1016/j.epsl.2007.08.028
- Burton MR, Oppenheimer C, Horrocks LA, Francis PW (2000) Remote sensing of CO₂ and H₂O emission rates from Masaya volcano, Nicaragua. *Geology* 28(10):915-918, doi: 10.1130/0091-7613(2000)28<915:RSO CAH>2.0.CO;2
- Cadle RD (1980) A comparison of volcanic with other fluxes of atmospheric trace gas constituents. *Rev Geophys Space Phys* 18:746-752
- Caliro S, Chiodini G, Galluzzo D, Granieri D, La Rocca M, Saccorotti G, Ventura G (2005) Recent activity of Nisyros volcano (Greece) inferred from structural, geochemical and seismological data. *Bull Volcanol* 67(4):358-369, doi: 10.1007/s00445-004-0381-7
- Camarda M, Gurrieri S, Valenza M (2006) CO₂ flux measurements in volcanic areas using the dynamic concentration method: Influence of soil permeability. *J Geophys Res* 111:B05202, doi: 10.1029/2005JB003898
- Carapezza ML, Federico C (2000) The contribution of fluid geochemistry to the volcano monitoring of Stromboli. *J Volcanol Geotherm Res* 95(1-4):227-245, doi: 10.1016/S0377-0273(99)00128-6
- Carapezza ML, Granieri D (2004) CO₂ soil flux at Vulcano (Italy): comparison between active and passive methods. *Appl Geochem* 19(1):73-88, doi: 10.1016/S0883-2927(03)00111-2
- Cardellini C, Chiodini G, Frondini F (2003) Application of stochastic simulation to CO₂ flux from soil: Mapping and quantification of gas release. *J Geophys Res Solid Earth* 108:2425, doi: 10.1029/2002JB002165
- Carn SA, Krotkov NA, Yang K, Krueger AJ (in press) Measuring global volcanic degassing with the Ozone Monitoring Instrument (OMI). *Spec Publ Geol Soc London*
- Cartigny P, Jendrzejewski N, Pineau F, Petit E, Javoy M (2001) Volatile (C, N, Ar) variability in MORB and the respective roles of mantle source heterogeneity and 25 degassing: the case study of the Southwest Indian ridge. *Earth Planet Sci Lett* 194:241-257
- Cheng ZH, Guo ZF, Zhang ML, Zhang LH (2012) CO₂ flux estimations of hot springs in the Tengchong Cenozoic volcanic field, Yunnan Province, SW China. *Acta Petrol Sinica* 28(4):1217-1224
- Chiodini G, Baldini A, Barberi F, Carapezza ML, Cardellini C, Frondini F, Granieri D, Ranaldi M (2007) Carbon dioxide degassing at Latera caldera (Italy): Evidence of geothermal reservoir and evaluation of its potential energy. *J Geophys Res Solid Earth* 112:B12204, doi: 10.1029/2006JB004896
- Chiodini G, Caliro S, Aiuppa A, Avino R, Granieri D, Moretti M, Parello F (2011) First ¹³C/¹²C isotopic characterisation of volcanic plume CO₂. *Bull Volcanol* 73:531-542, doi: 10.1007/s00445-010-0423-2

- Chiodini G, Cardellini C, Amato A, Boschi E, Caliro S, Frondini F, Ventura G (2004) Carbon dioxide Earth degassing and seismogenesis in central and southern Italy. *Geophys Res Lett* 31(7):L07615, doi: 10.1029/2004GL019480
- Chiodini G, Cioni R, Guidi M, Raco B, Marini L (1998) Soil CO₂ flux measurements in volcanic and geothermal areas. *Appl Geochem* 13:543-552
- Chiodini G, Frondini F (2001) Carbon dioxide degassing from the Albani Hills volcanic region, Central Italy. *Chem Geol* 177(1-2):67-83, doi: 10.1016/S0009-2541(00)00382-X
- Chiodini G, Frondini F, Cardellini C, Granieri D, Marini L, Ventura G (2001) CO₂ degassing and energy release at Solfatara volcano, Campi Flegrei, Italy. *J Geophys Res Solid Earth* 106:16,213-16,221, doi: 10.1029/2001JB000246
- Chiodini G, Frondini F, Kerrick DM, Rogie J, Parello F, Peruzzi L, Zanzari AR (1999) Quantification of deep CO₂ fluxes from Central Italy. Examples of carbon balance for regional aquifers and of soil diffuse degassing. *Chem Geol* 159(1-4):205-222, doi: 10.1016/S0009-2541(99)00030-3
- Chiodini G, Frondini F, Raco B (1996) Diffuse emission of CO₂ from the Fossa crater, Vulcano Island (Italy). *Bull Volcanol* 58:41-50
- Chiodini G, Granieri D, Avino R, Caliro S, Costa A, Minopoli C, Vilardo G (2010) Non-volcanic CO₂ Earth degassing: Case of Mefite d'Ansanto (southern Apennines), Italy. *Geophys Res Lett* 37:L11303, doi: 10.1029/2010GL042858
- Chiodini G, Granieri D, Avino R, Caliro S, Costa A, Werner C (2005) Carbon dioxide diffuse degassing and estimation of heat release from volcanic and hydrothermal systems, *J Geophys Res Solid Earth* 110:B08204, doi: 10.1029/2004JB003542
- Corliss JB, Dymond J, Forgon LI, Edmond JM, von Herzen RP, Ballard RD, Green K, Williams D, Bainbridge A, Crane K, van Andel TH (1979) Submarine thermal springs on the Galápagos Rift. *Science* 203:1073-1083
- Crispa D, Atlasb RM, Breonc F-M, Browna LR, Burrowsd JP, Ciaise P, Connore BJ, Doneyf SC, Fungg IY, Jacobh DJ, Milleri CE, O'Brienj D, Pawsonb S, Randersonk JT, Raynerj P, Salawitcha RJ, Sandera SP, Sena B, Stephensm GL, Tansn PP, Toona GC, Wennbergk PO, Wofsyh SC, Yungk YL, Kuangk Z, Chudasamaa B, Spraguea G, Weissa B, Pollocko R, Kenyonp D, Schrollp S (2004) The orbiting carbon observatory (OCO) mission. Trace Constituents in the Troposphere and Lower Stratosphere: *Adv Space Res* 34(4), doi: 10.1016/j.asr.2003.08.062
- Crisp JA (1984) Rates of magma emplacement and volcanic output. *J Volcanol Geotherm Res* 20(3-4):177-211, doi: 10.1016/0377-0273(84)90039-8
- Crossey LJ, Karlstrom KE, Springer AE, Newell D, Hilton DR, Fischer T (2009) Degassing of mantle-derived CO₂ and He from springs in the southern Colorado Plateau region-Neotectonic connections and implications for groundwater systems. *Geol Soc Am Bull* 121(7-8):1034-1053, doi: 10.1130/B26394.1
- Crutzen PJ (1976) The possible importance of COS for the sulfate layer of the stratosphere. *Geophys Res Lett* 3:73-76
- Cruz JV, Coutinho RM, Carvalho MR, Oskarsson N, Gislason SR (1999) Chemistry of waters from Furnas volcano, São Miguel, Azores: fluxes of volcanic carbon dioxide and leached material. *J Volcanol Geotherm Res* 92:151-167
- D'Alessandro W, Giammanco S, Parello F, Valenza M (1997) CO₂ output and $\delta^{13}\text{C}(\text{CO}_2)$ from Mount Etna as indicators of degassing of shallow asthenosphere. *Bull Volcanol* 58:455-458
- Dasgupta R, Hirschmann MM (2010) The deep carbon cycle and melting in Earth's interior. *Earth Planet Sci Lett (Frontiers)* 298:1-13
- Delgado H, Piedad-Sanchez N, Galvan L, Julio T, Alvarez M, Cardenas L (1998) CO₂ flux measurements at Popocatepetl volcano: II Magnitude of emissions and significance. *EOS Trans/Supplement* 79, no 10, p F926
- Des Marais DJ, Moore JG (1984) Carbon and its isotopes in mid-oceanic basaltic glasses. *Earth Planet Sci Lett* 69:43-57, doi: 10.1016/0012-821X(84)90073-6
- Diaz JA, Pieri D, Arkin CR, Gore E, Griffin TP, Fladeland M, Bland G, Soto C, Madrigal Y, Castillo D, Rojas E, Achi S (2010) Utilization of in situ airborne MS-based instrumentation for the study of gaseous emissions at active volcanoes. *Int J Mass Spectrom* 295(3):105-112, doi: 10.1016/j.ijms.2010.04.013
- Dixon JE, Clague DA (2001) Volatiles in basaltic glasses from Loihi seamount, Hawaii: evidence for a relatively dry plume component. *J Petrol* 42:627-654
- Doukas MP, McGee KA (2007) A compilation of gas emission-rate data from volcanoes of Cook Inlet (Spurr, Crater Peak, Redoubt, Iliamna, and Augustine) and Alaska Peninsula (Douglas, Fourpeaked, Griggs, Mageik, Martin, Peulik, Ukinrek Maars, and Veniaminof), Alaska, from 1995-2006. *US Geol Soc Open-File Report* 2007-1400
- Edmonds M, Aiuppa A, Humphreys M, Moretti R, Giudice G, Martin RS, Herd RA, Christopher T (2010) Excess volatiles supplied by mingling of mafic magma at an andesite arc volcano. *Geochem Geophys Geosyst* 11:Q04005, doi: 10.1029/2009GC002781

- Edmonds M, Herd RA, Galle B, Oppenheimer CM (2003) Automated, high time-resolution measurements of SO₂ flux at Soufriere Hills Volcano, Montserrat. *Bull Volcanol* 65:578-586, doi: 10.1007/s00445-003-0286-x
- Eichelberger JC, Carrigan CR, Westrich HR, Shannon JR (1986) Non-explosive volcanism. *Nature* 323:598-602, doi: 10.1038/323598a0
- Elderfield H, Schultz A (1996) Mid-ocean ridge hydrothermal fluxes and the chemical composition of the ocean. *Annu Rev Earth Planet Sci* 24:191-224
- Etiopio G, Beneduce P, Calcara M, Favali P, Frugoni F, Schiattarella M, Smriglio G (1999) Structural pattern and CO₂-CH₄ degassing of Ustica Island, Southern Tyrrhenian basin. *J Volcanol Geotherm Res* 88:291-304
- Etiopio G, Milkov AV, Derbyshire E (2008) Did geologic emissions of methane play any role in Quaternary climate change? *Global Planet Change* 61(1-2):79-88, doi:10.1016/j.gloplacha.2007.08.008
- Evans WC, Bergfeld D, McGimsey RG, Hunt AG (2009) Diffuse gas emissions at the Ukinrek Maars, Alaska: Implications for magmatic degassing and volcanic monitoring. *Appl Geochem* 24(4):527-535, doi: 10.1016/j.apgeochem.2008.12.007
- Farley KA, Maier-Reimer E, Schlosser P, Broecker WS (1995). Constraints on mantle ³He fluxes and deep-sea circulation from an oceanic general circulation model. *J Geophys Res* 100:3829-3839, doi: 10.1029/94JB02913
- Favara R, Giammanco S, Inguaggiato S, Pecoraino G (2001) Preliminary estimate of CO₂ output from Pantelleria Island volcano (Sicily, Italy): evidence of active mantle degassing. *Appl Geochem* 16:883-894
- Feulner G (2012) The faint young Sun problem. *Rev Geophys* 50:RG2006, doi: 10.1029/2011RG000375
- Fischer TP (2008) Fluxes of volatiles (H₂O, CO₂, N₂, Cl, F) from arc volcanoes. *Geochem J* 42(1):21-38
- Fischer TP, Giggenbach WF, Sano Y, Williams SN (1998) Fluxes and sources of volatiles discharged from Kudryavy, a subduction zone volcano, Kurile Islands. *Earth Planet Sci Lett* 160(1-2):81-96, doi: 10.1016/S0012-821X(98)00086-7
- Fischer TP, Marty B (2005) Volatile abundances in the sub-arc mantle: insights from volcanic and hydrothermal gas discharges. *J Volcanol Geotherm Res* 140:205-216
- Francis PW, Oppenheimer C, Stevenson D (1993) Endogenous growth of persistently active volcanoes. *Nature* 366:554-557, doi: 10.1038/366554a0
- Friedlingstein P, Houghton RA, Marland G, Hackler J, Boden TA, Conway TJ, Canadell JG, Raupach MR, Ciais P, Le Quééré C (2010) Update on CO₂ emissions. *Nat Geosci* 3(12):811-812, doi: 10.1038/ngeo1022
- Fronzini F, Chiodini G, Caliro S, Cardellini C, Granieri D, Ventura G (2004) Diffuse CO₂ degassing at Vesuvio, Italy. *Bull Volcanol* 66(7):642-651, doi: 10.1007/s00445-004-0346-x
- Füri ED, Hilton R, Tryon MD, Brown KM, McMurtry GM, Brückmann W, Wheat CG (2010) Carbon release from submarine seeps at the Costa Rica fore-arc: Implications for the volatile cycle at the Central America convergent margin. *Geochem Geophys Geosyst* 11:Q04S21, doi: 10.1029/2009GC002810
- Gaillardet J, Dupre B, Louvat P, Allegre CJ (1999) Global silicate weathering and CO₂ consumption rates deduced from the chemistry of the large rivers. *Chem Geol* 159(1-4):3-30
- Galle B, Johansson M, Rivera C, Zhang Y, Kihlman M, Kern C, Lehmann T, Platt U, Arellano S, Hidalgo S (2010) Network for Observation of Volcanic and Atmospheric Change (NOVAC)—A global network for volcanic gas monitoring: Network layout and instrument description. *J Geophys Res* 115:D05304, doi: 10.1029/2009JD011823
- Gerlach TM (1989) Degassing of carbon dioxide from basaltic magma at spreading centers: II. Mid-oceanic ridge basalts. *J Volcanol Geotherm Res* 39:231-232
- Gerlach TM (1991) Present-day CO₂ emissions from volcanoes. *EOS Trans, Am Geophys Union* 72 (23):249-255
- Gerlach TM (2011) Volcanic versus anthropogenic carbon dioxide. *EOS Trans*, 92(24):201-208
- Gerlach TM, Delgado H, McGee KA, Doukas MP, Venegas JJ, Cardenas L (1997) Application of the LI-COR CO₂ analyzer to volcanic plumes: A case study, volcan Popocatepetl, Mexico, June 7 and 10, 1995. *J Geophys Res Solid Earth* 102:8005-8019, doi: 10.1029/96JB03887
- Gerlach TM, McGee KA, Elias T, Sutton AJ, Doukas MP (2002) Carbon dioxide emission rate of Kilauea Volcano: Implications for primary magma and the summit reservoir. *J Geophys Res Solid Earth* 107:2189, doi: 10.1029/2001JB000407
- Gerlach TM, McGee KA, Sutton AJ, Elias T (1998) Rates of volcanic CO₂ degassing from airborne determinations of SO₂, emission rates and plume CO₂/SO₂: Test study at Pu'u 'O'o cone, Kilauea volcano, Hawaii. *Geophys Res Lett* 25(14):2675-2678, doi: 10.1029/98GL02030
- Giammanco S, Parello F, Gambardella B, Schifano R, Pizzu S, Galante G (2007) Focused and diffuse effluxes of CO₂ from mud volcanoes and mofettes south of Mt. Etna (Italy). *J Volcanol Geotherm Res* 165:46-63, doi: 10.1016/j.jvolgeores.2007.04.010
- Giggenbach WF, Goguel RL (1989) Methods for the collection and analysis of geothermal and volcanic water and gas samples. Department of Scientific and Industrial Research, Chemistry Division, Report 2401

- Goff F, Love SP, Warren RG, Counce D, Obenholzner J, Siebe C, Schmidt SC (2001) Passive infrared remote sensing evidence for large, intermittent CO₂ emissions at Popocatepetl volcano, Mexico. *Chem Geol* 177(1-2):133-156, doi: 10.1016/S0009-2541(00)00387-9
- Graham D, Sarda P (1991) Reply to comment by TM Gerlach on "Mid-ocean ridge popping rocks: implications for degassing at ridge crests". *Earth Planet Sci Lett* 105: 568-573
- Greenland LP, Rose WI, Stokes JB (1985) An estimate of gas emissions and magmatic gas content from Kilauea volcano. *Geochim Cosmochim Acta* 49(1):125-129, doi: 10.1016/0016-7037(85)90196-6
- Hager SA, Gerlach TM, Wallace PJ (2008) Summit CO₂ emission rates by the CO₂/SO₂ ratio method at Kilauea Volcano, Hawai'i, during a period of sustained inflation. *J Volcanol Geotherm Res* 177(4):875-882, doi: 10.1016/j.jvolgeores.2008.06.033
- Halmer MM, Schmincke HU, Graf HF (2002) The annual volcanic gas input into the atmosphere, in particular into the stratosphere: A global data set for the past 100 years. *J Volcanol Geotherm Res* 115:511-528
- Hernández P, Pérez N, Salazar J, Sato M, Notsu K, Wakita H (2000) Soil gas CO₂, CH₄, and H₂ distribution in and around Las Cañadas caldera, Tenerife, Canary Islands, Spain. *J Volcanol Geotherm Res* 103:425-438
- Hernández PA, Pérez NM, Fridriksson T, Egbert J, Ilyinskaya E, Thárhallsson A, Ívarsson G, Gíslason G, Gunnarsson I, Jónsson B, Padrón E, Melián G, Mori T, Notsu K (2012) Diffuse volcanic degassing and thermal energy release from Hengill volcanic system, Iceland. *Bull Volcanol* 74:2435-2448, doi: 10.1007/s00445-012-0673-2
- Hernández PA, Notsu K, Okada H, Mori T, Sato M, Barahona F, Pérez NM (2006) Diffuse emission of CO₂ from Showa-Shinzan, Hokkaido, Japan: A sign of volcanic dome degassing. *Pure Appl Geophys* 163(4):869-881, doi: 10.1007/s00024-006-0038-x
- Hernández PA, Notsu K, Salazar JM, Mori T, Natale G, Okada H, Virgili G, Shimoike Y, Sato M, Pérez NM (2001b) Carbon dioxide degassing by advective flow from Usu volcano, Japan. *Science* 292:83-86
- Hernández PA, Notsu K, Tsurumi M, Mori T, Ohno M, Shimoike Y, Salazar J, Pérez NM (2003) Carbon dioxide emissions from soils at Hakkoda, north Japan. *J Geophys Res* 108:2210, doi: 10.1029/2002JB001847
- Hernández PA, Salazar JM, Shimoike Y, Mori T, Notsu K, Pérez NM (2001a) Diffuse emission of CO₂ from Miyakejima volcano. *Japan Chem Geol* 177:175-185
- Hilton DR, Fischer TP, Marty B (2002) Noble gases and volatile recycling at subduction zones. *Rev Mineral Geochem* 47:319-370
- Holloway JR (1998) Graphite-melt equilibria during mantle melting: constraints on CO₂ in MORB magmas and the carbon content of the mantle. *Chem Geol* 147:89-97
- Hornafius JS, Quigley D, Luyendyk BP (1999) The world's most spectacular marine hydrocarbon seeps (Coal Oil Point, Santa Barbara Channel, California): quantification of emissions. *J Geophys Res* 104:20703-20711
- Inguaggiato S, Mazot A, Diliberto IS, Inguaggiato C, Madonia P, Rouwet D, Vita F (2012) Total CO₂ output from Vulcano island (Aeolian Islands, Italy). *Geochem Geophys Geosyst* 13:Q02012, doi: 10.1029/2011GC003920
- Irwin WP, Barnes I (1980) Tectonic relations of carbon dioxide discharges and earthquakes. *J Geophys Res* 85:3115-3121, doi: 10.1029/JB085iB06p03115
- Italiano F, Martelli M, Martinelli G, Nuccio PM (2000) Geochemical evidence of melt intrusions along lithospheric faults of the Southern Apennines, Italy: geodynamic and seismogenic implications. *J Geophys Res* 105:13569-13578
- James ER, Manga M, Rose TP (1999) CO₂ degassing in the Oregon cascades. *Geology* 27(9):823-826, doi: 10.1130/0091-7613(1999)027<0823:CDITOC>2.3.CO;2
- Jaupart C, Vergnolle S (1988) Laboratory models of Hawaiian and Strombolian eruptions. *Nature* 331:58-60, doi: 10.1038/331058a0
- Javoy M, Pineau F (1991) The volatiles record of a bopping rock from the Mid-Atlantic ridge at 148N: chemical and isotopic composition of a gas trapped in the vesicles. *Earth Planet Sci Lett* 107:598-611
- Kanemasu ET, Powers WL, Sij JW (1974) Field chamber measurements of CO₂ flux from soil surface. *Soil Sci* 118(4):233-237, doi: 10.1097/00010694-197410000-00001
- Kazahaya K, Shinohara H, Saito G (1994) Excessive degassing of Izu-Oshima volcano: Magma convection in a conduit. *Bull Volcanol* 56:207-216, doi: 10.1007/BF00279605
- Kelley DS, Lilley MD, Fruh-Green GL (2004) Volatiles in submarine environments: Food for life. *In: The Subseafloor Biosphere at Mid-Ocean Ridges*. *Geophys Monogr Ser* 144, Wilcock W, DeLong EF, Kelley DS, Baross JA, Craig Cary S (eds) AGU, Washington, D.C., p 167-190
- Kern C, Deutschmann T, Werner C, Sutton AJ, Elias T, Kelly PJ (2012) Improving the accuracy of SO₂ column densities and emission rates obtained from upward-looking UV-spectroscopic measurements of volcanic plumes by taking realistic radiative transfer into account. *J Geophys Res* 117:D20302, doi: 10.1029/2012JD017936
- Kerrick DM (2001) Present and past non-anthropogenic CO₂ degassing from the solid earth. *Rev Geophys* 39(4):565-585

- Kerrick DM, McKibben MA, Seward TM, Caldeira K (1995) Convective hydrothermal CO₂ emission from high heat flow regions. *Chem Geol* 121:285-293
- Koepnick KW, Brantley SL, Thompson JM, Rowe GL, Nyblade AA, Moshy C (1996) Volatile emissions from the crater and flank of Oldoinyo Lengai, Tanzania. *J Geophys Res* 101:13819-13830
- Kucera CL, Kirkham DR (1971) Soil respiration studies in tall grass prairie in Missouri. *Ecology* 52(5):912-915, doi: 10.2307/1936043
- La Spina A, Burton M, Harig R, Murè F, Rausch P, Jordan M, Caltabiano T (2013) New insights into volcanic processes at Stromboli from Cerberus, a remote-controlled open-path FTIR scanner. *J Volcanol Geotherm Res* 249:66-76
- La Spina A, Burton M, Salerno GG (2010) Unravelling the processes controlling gas emissions from the central and northeast craters of Mt. Etna. *J Volcanol Geotherm Res* 198(3-4):368-376
- Le Guern F (1987) Mechanism of energy transfer in the lava lake of Niragongo (Zaire), 1959-1977. *J Volcanol Geotherm Res* 31(1-2):17-31, doi: 10.1016/0377-0273(87)90003-5
- Le Guern F, Carbonnelle J, Taziéff H (1979) Erta 'Ale lava lake: heat and gas transfer to the atmosphere. *J Volcanol Geotherm Res* 6:27-48
- Lewicki JL, Fischer ML, Hilley GE (2008) Six-week time series of eddy covariance CO₂ flux at Mammoth Mountain, California: Performance evaluation and role of meteorological forcing. *J Volcanol Geotherm Res* 171(3-4):178-190, doi: 10.1016/j.jvolgeores.2007.11.029
- Lopez T, Ushakov S, Izbekov P, Tassi F, Cahill C, Neill O, Werner C (in press) Constraints on magma processes, subsurface conditions, and total volatile flux at Bezymianny Volcano in 2007-2010 from direct and remote volcanic gas measurements. *J Volcanol Geotherm Res*, doi: 10.1016/j.jvolgeores.2012.10.015
- Lupton J, Lilliey M, Butterfield D, Evans L, Embley R, Massoth G, Christenson B, Nakamura K, Schmidt M (2008) Venting of a separate CO₂-rich gas phase from submarine arc volcanoes: Examples from the Mariana and Tonga-Kermadec arcs. *J Geophys Res* 113:B08S12, doi: 10.1029/2007JB005467
- Marrero R, López DL, Hernández PA, Pérez NM (2008) Carbon dioxide discharged through the Las Cañadas Aquifer, Tenerife, Canary Islands. *Pure Appl Geophys* 165:147-172, doi: 10.1007/s00024-007-0287-3
- Martin RS, Sawyer GM, Spampinato L, Salerno GG, Ramirez C, Ilyinskaya E, Witt MLI, Mather TA, Watson IM, Phillips JC, Oppenheimer C (2010) A total volatile inventory for Masaya Volcano, Nicaragua. *J Geophys Res Solid Earth* 115:B09215, doi: 10.1029/2010JB007480
- Marty B, Jambon A (1987) C³He in volatile fluxes from the solid earth: implications for carbon geodynamics. *Earth Planet Sci Lett* 83:16-26
- Marty B, Le Cloarec MF (1992) Helium-3 and CO₂ fluxes from subaerial volcanoes estimated from polonium-210 emissions. *J Volcanol Geotherm Res* 53:67-72
- Marty B, Tolstikhin IN (1998) CO₂ fluxes from mid-ocean ridges, arcs and plumes. *Chem Geol* 145:233-248
- Marty B, Zimmerman L (1999) Volatiles (He, C, N, Ar) in mid-ocean ridge basalts: assessment of shallow-level fractionation and characterization of source composition. *Geochim Cosmochim Acta* 63(21):3619-3633, doi: 10.1016/S0016-7037(99)00169-6
- Massoth GJ, Milburn HB, Hammond SR, Butterfield DA, McDuff RE, Lupton JE (1989) The geochemistry of submarine venting fluids at Axial Volcano, Juan de Fuca Ridge: New sampling methods and a VENTS program rationale. In: *Global Venting, Midwater, and Benthic Ecological Processes*. National Undersea Research Program, Research Report 88-4. De Luca MP, Babb I (eds) US Department of Commerce: National Oceanic and Atmospheric Administration, p 29-59
- Matthews SJ, Jones AP, Bristow CS (1992) A simple magma-mixing model for sulfur behavior in calc-alkaline volcanic rocks: mineralogical evidence from Mount Pinatubo 1991 eruption. *J Geol Soc London* 149: 863-866
- Mazot A, Taran Y (2009) CO₂ flux from the volcanic lake of El Chichón (Mexico): *Geofis Int* 48(1):73-83
- McGee KA, Doukas MP, Gerlach TM (2001) Quiescent hydrogen sulfide and carbon dioxide degassing from Mount Baker, Washington. *Geophys Res Lett* 28(23):4479-4482, doi: 10.1029/2001GL013250
- McGonigle AJS, Aiuppa A, Giudice G, Tamburello G, Hodson AJ, Gurrieri S (2008) Unmanned aerial vehicle measurements of volcanic carbon dioxide fluxes. *Geophys Res Lett* 35(6):L06303, doi: 10.1029/2007GL032508
- Menand T, Phillips HC (2007) Gas segregation in dykes and sills. *J Volcanol Geotherm Res* 159(4):393-408, doi: 10.1016/j.jvolgeores.2006.08.003
- Merucci L, Burton M, Corradini S, Salerno GG (2011) Reconstruction of SO₂ flux emission chronology from space-based measurements. *J Volcanol Geotherm Res* 206(3-4):80-87, doi: 10.1016/j.jvolgeores.2011.07.002
- Métrich N, Allard P, Aiuppa A, Bani P, Bertagnini A, Shinohara H, Parello F, Di Muro A, Garaebiti E, Belhadj O, Massare D (2011) Magma and volatile supply to post-collapse volcanism and block resurgence in Siwi Caldera (Tanna Island, Vanuatu Arc). *J Petrol* 52(6):1077-1105, doi: 10.1093/petrology/egr019
- Mori T, Burton MR (2006) The SO₂ camera: A simple, fast and cheap method for ground-based imaging of SO₂ in volcanic plumes. *Geophys Res Lett* 33(24):L24804, doi: 10.1029/2006GL027916
- Mori T, Hernández PA, Salazar JML, Pérez NM, Notsu K (2001) An in situ method for measuring CO₂ flux from volcanic-hydrothermal fumaroles. *Chem Geol* 177(1-2):85-99, doi: 10.1016/S0009-2541(00)00384-3

- Mori T, Notsu K (1997) Remote CO, COS, CO₂, SO₂, HCl detection and temperature estimation of volcanic gas. *Geophys Res Lett* 24(16):2047-2050, doi: 10.1029/97GL52058
- Mörner NA, Etiope G (2002) Carbon degassing from the lithosphere. *Global Planet Change* 33:185-203
- Naughton JJ, Derby JV, Glover RB (1969) Infrared measurements on volcanic gas and fume: Kilauea eruption, 1968. *J Geophys Res* 74:3273-3277
- Oppenheimer C, Kyle PR (2008) Probing the magma plumbing of Erebus volcano, Antarctica, by open-path FTIR spectroscopy of gas emissions. *J Volcanol Geotherm Res* 177(3):743-754, doi: 10.1016/j.jvolgeores.2007.08.022
- Oppenheimer C, Scaillet B, Martin RS (2011) Sulfur degassing from volcanoes: source conditions, surveillance, plume chemistry and earth system impacts. *In: Sulfur in magmas and melts: its importance for natural and technical processes. Rev Mineral Geochem* 73:363-422, doi: 10.2138/rmg.2011.73.13
- Óskarsson F, Fridriksson T (2011) Reykjanes production field. *Geochemical monitoring in 2010. ÍSOR Report* 2011/050, Iceland GeoSurvey, Reykjavík, p 51
- Padron E, Hernández PA, Pérez NM, Toulkeridis T, Melian G, Barrancos J, Virgili G, Sumino H, Notsu K (2012) Fumarole/plume and diffuse CO₂ emission from Sierra Negra caldera, Galapagos archipelago. *Bull Volcanol* 74(6):1509-1519, doi: 10.1007/s00445-012-0610-4
- Pallister JS, Hoblitt RP, Reyes AG (1992) A basalt trigger for the 1991 eruptions of Pinatubo volcano. *Nature* 356(6368):426-428, doi: 10.1038/356426a0
- Parkinson KJ (1981) An improved method for measuring soil respiration in the field. *J Appl Ecol* 18:221-228
- Pecoraino G, Brusca L, D'Alessandro W, Giammanco S, Inguaggiato S, Longo M (2005) Total CO₂ output from Ischia Island volcano (Italy). *Geochim J* 39:451-458
- Pérez N, Melian G, Salazar J, Saballos A, Alvarez J, Segura F, Hernández P, Notsu K (2000) Diffuse degassing of CO₂ from Masaya Caldera, Nicaragua, Central America. *EOS Trans, Am Geophys Union* 81(48), Fall Meet Suppl, 2000
- Pérez NM et al. (2011) Global CO₂ emission from volcanic lakes. *Geology* 39:235-238, doi: 10.1130/G31586.1
- Pérez NM, Nakai S, Wakita H, Notsu K, Talai B (1998) Anomalous diffuse degassing of Helium-3 and CO₂ related to the active ring-fault structure at Rabaul Caldera, Papua New Guinea. *Abstract AGU Fall Meet*, V12B-11
- Pili E, Kennedy BM, Conrad ME, Gratier J-P (2011) Isotopic evidence for the infiltration of mantle and metamorphic CO₂-H₂O fluids from below in faulted rocks from the San Andreas Fault system. *Chem Geol* 281(3-4):242-252, doi: 10.1016/j.chemgeo.2010.12.011
- Reiners WA (1968) Carbon dioxide evolution from the floor of three Minnesota forests. *Ecology* 49:471-483
- Resing JA, Lupton JE, Feely RA, Lilley MD (2004) CO₂ and ³He in hydrothermal plumes: implications for mid-ocean ridge CO₂ flux. *Earth Planet Sci Lett* 226:449-464
- Roberts TJ, Braban CF, Oppenheimer C, Martin RS, Freshwater RA, Dawson DH, Griffiths PT, Cox RA, Saffell JR, Jones RL (2012) Electrochemical sensing of volcanic gases, *Chem Geol* 332-333:74-91, doi: 10.1016/j.chemgeo.2012.08.027
- Rogie JD, Kerrick DM, Chiodini G, Frondini F (2000) Flux measurements of nonvolcanic CO₂ emission from some vents in central Italy. *J Geophys Res Solid Earth* 105:8425-8445, doi: 10.1029/1999JB900430
- Rose TP, Davison ML (1996) Radiocarbon in hydrologic systems containing dissolved magmatic carbon dioxide. *Science* 273(5280):1367-1370, doi: 10.1126/science.273.5280.1367
- Rose WI, Chuan RL, Giggenbach WF, Kyle PR, Symonds RB (1986) Rates of sulfur dioxide and particle emissions from White Island volcano, New Zealand, and an estimate of the total flux of major gaseous species. *Bull Volcanol* 48(4), doi: 181-188
- Saal AE, Hauri EH, Langmuir CH, Perfit MR (2002) Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of earth's upper mantle. *Nature* 419:451-455
- Salazar JML, Hernández PA, Pérez NM, Melian G, Alvarez J, Segura F, Notsu K (2001) Diffuse emission of carbon dioxide from Cerro Negro volcano, Nicaragua, Central America. *Geophys Res Lett* 28(22):4275-4278, doi: 10.1029/2001GL013709
- Salazar JML, Hernández PA, Pérez NM, Olmos R, Barahona F, Cartagena R, Soriano T, Lopez DL, Sumino H, Notsu K (2004) Spatial and temporal variations of diffuse CO₂ degassing at the Santa Ana-Izalco-Coatepeque volcanic complex, El Salvador, Central America. *Geol Soc Am Special Papers* 375:135-146, doi: 10.1130/0-8137-2375-2.135
- Sano Y, Williams SN (1996) Fluxes of mantle and subducted carbon along convergent plate boundaries. *Geophys Res Lett* 23(20):2749-2752
- Sansone FJ, Mottl MJ, Olson EJ, Wheat CG, Lilley MD (1998) CO₂-depleted fluids from mid-ocean ridge-flank hydrothermal springs. *Geochim Cosmochim Acta* 62(13):2247-2252, doi: 10.1016/S0016-7037(98)00135-5
- Sarda SP, Graham D (1990) Mid-ocean ridge popping rocks: implications for degassing at ridge crests. *Earth Planet Sci Lett* 97:268-289

- Sawyer GM, Carn SA, Tsanev VI, Oppenheimer C, Burton MR (2008a) Investigation into magma degassing at Nyiragongo volcano, Democratic Republic of the Congo. *Geochem Geophys Geosyst* 9:Q02017, doi: 10.1029/2007GC001829
- Sawyer GM, Oppenheimer C, Tsanev VI, Yirgu G (2008b) Magmatic degassing at Erta 'Ale volcano, Ethiopia. *J Volcanol Geotherm Res* 178:837-846, doi: 10.1016/j.jvolgeores.2008.09.017
- Sawyer GM, Salerno GG, Le Blond J, Martin RS, Spampinato L, Roberts T, Tsanev VI, Mather TA, Witt MLI, Oppenheimer C (2011) Gas and particle emissions from Villarrica Volcano, Chile. *J Volcanol Geotherm Res* 203(1-2):62-75, doi: 10.1016/j.jvolgeores.2011.04.003
- Self S (2006) The effects and consequences of very large explosive eruptions. *Philos Trans R Soc A* 364:2073-2097, doi: 10.1098/rsta.2006.1814
- Seward TM, Kerrick DM (1996) Hydrothermal CO₂ emission from the Taupo Volcanic Zone, New Zealand. *Earth Planet Sci Lett* 139(1-2):105-113, doi: 10.1016/0012-821X(96)00011-8
- Shimoike Y, Kazahaya K, Shinohara H (2002) Soil gas emission of volcanic CO₂ at Satsuma-Iwojima volcano, Japan. *Earth Planet Sci Lett* 54(3):239-247
- Shinohara H (2005) A new technique to estimate volcanic gas composition: plume measurements with a portable multi-sensor system. *J Volcanol Geotherm Res* 143(4):319-333
- Shinohara H, Kazahaya K, Saito G, Fukui K, Odai M (2003) Variation of CO₂/SO₂ ratio in volcanic plumes of Miyakejima: Stable degassing deduced from heliborne measurements. *Geophys Res Lett* 30(5):1208, doi: 10.1029/2002GL016105
- Siebert L, Simkin T (2002) *Volcanoes of the world: an illustrated catalogue of Holocene volcanoes and their eruptions*. Smithsonian Institution, Global Volcanism Program Digital Information Series, GVP-3, (<http://www.volcano.si.edu/world/>)
- Smith TEL, Wooster MJ, Tattaris M, Griffith DWT (2011) Absolute accuracy and sensitivity analysis of OP-FTIR retrievals of CO₂, CH₄ and CO over concentrations representative of "clean air" and "polluted plumes". *Atmos Meas Tech* 4:97-116, doi: 10.5194/amt-4-97-2011
- Sorey ML, Evans WC, Kennedy BM, Farrar CD, Hainsworth LJ, Hausback B (1998) Carbon dioxide and helium emissions from a reservoir of magmatic gas beneath Mammoth Mountain, California. *J Geophys Res Solid Earth* 103:15,503-15,323, doi: 10.1029/98JB01389
- Spilliaert N, Allard P, Métrich N, Sobolev AV (2006) Melt inclusion record of the conditions of ascent, degassing, and extrusion of volatile-rich alkali basalt during the powerful 2002 flank eruption of Mount Etna (Italy). *J Geophys Res* 111:B04203, doi: 10.1029/2005JB003934
- Spinetti C, Carrere V, Buongiorno MF, Sutton AJ, Elias T (2008) Carbon dioxide of Pu'u'Ō'o volcanic plume at Kilauea retrieved by AVIRIS hyperspectral data. *Remote Sens Environ* 112(6):3192-3199, doi: 10.1016/j.rse.2008.03.010
- Symonds RB, Reed MH, Rose WI (1992) Origin, speciation, and fluxes of trace-element gases at Augustine volcano, Alaska – Insights into magma degassing and fumarolic processes. *Geochim Cosmochim Acta* 56(2):633-657, doi: 10.1016/0016-7037(92)90087-Y
- Tonani F, Miele G (1991) Methods for measuring flow of carbon dioxide through soil in volcanic setting. *In: International Conference on Active Volcanoes and Risk Mitigation, IAVCEI, Naples, Italy, 27 Aug. to 1 Sept.*
- Toutain J-P, Sortino F, Baubron J-C, Richon P, Surono, Sumarti S, Nonell A (2009) Structure and CO₂ budget of Merapi volcano during inter-eruptive periods. *Bull Volcanol* 71(7):815-826, doi: 10.1007/s00445-009-0266-x
- Troll VR, Hilton DR, Jolis EM, Chadwick JP, Blythe LS, Deegan FM, Schwarzkopf LM, Zimmer M (2012) Crustal CO₂ liberation during the 2006 eruption and earthquake events at Merapi volcano, Indonesia. *Geophys Res Lett* 39:L11302, doi: 10.1029/2012GL051307
- Varekamp JC, Kreulen R, Poorter RPE, Van Bergen MJ (1992) Carbon sources in arc volcanism, with implications for the carbon cycle. *Terra Nova* 4(3):363-373, doi: 10.1111/j.1365-3121.1992.tb00825.x
- Varley NR, Armienta MA (2001) The absence of diffuse degassing at Popocatepetl volcano, Mexico. *Chem Geol* 177:157-173
- Von Damm KL, Edmond JM, Grant G, Measures CI, Walden B, Weiss RF (1985) Chemistry of submarine hydrothermal solutions at 21°N, East Pacific Rise. *Geochim Cosmochim Acta* 49:2197-2220
- Walker JCG, Hays PB, Kasting JF (1981) A negative feedback mechanism for the long-term stabilization of Earth's surface temperature. *J Geophys Res* 86:9776-9782, doi: 10.1029/JC086iC10p09776
- Wallace PJ (2005) Volatiles in subduction zone magmas: concentrations and fluxes based on melt inclusion and volcanic gas data. *J Volcanol Geotherm Res* 140(1-3):217-240, doi: 10.1016/j.jvolgeores.2004.07.023
- Wallace PJ, Gerlach TM (1994) Magmatic vapor source for sulfur-dioxide released during volcanic-eruptions - evidence from Mount Pinatubo. *Science* 265(5171):497-499
- Wang WC, Yung YL, Lacis AA, Mo T, Hansen JE (1976) Greenhouse effects due to man-made perturbations of trace gases. *Science* 194(4266): 685-690

- Wardell LJ, Kyle PR, Campbell AR (2003) Carbon dioxide emissions from fumarolic ice towers, Mount Erebus volcano, Antarctica. *In: Volcanic Degassing*. Oppenheimer C, Pyle DM, Barclay J (eds) Geol Soc London, Special Publications 213:213-246
- Wardell LJ, Kyle PR, Chaffin C (2004) Carbon dioxide and carbon monoxide emission rates from an alkaline intra-plate volcano: Mt. Erebus, Antarctica. *J Volcanol Geotherm Res* 131(1-2):109-121, doi: 10.1016/S0377-0273(03)00320-2
- Wardell LJ, Kyle PR, Dunbar N, Christenson B (2001) White Island volcano, New Zealand: carbon dioxide and sulfur dioxide emission rates and melt inclusion studies. *Chem Geol* 177(1-2):187-200, doi: 10.1016/S0009-2541(00)00391-0
- Watts SF (2000) The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide. *Atmos Environ* 34:761-779
- Werner C, Brantley S (2003) CO₂ emissions from the Yellowstone volcanic system. *Geochem Geophys Geosyst* 4:1061, doi: 10.1029/2002GC000473
- Werner C, Evans WC, Poland M, Tucker DS, Doukas MP (2009) Long-term changes in quiescent degassing at Mount Baker Volcano, Washington, USA; Evidence for a stalled intrusion in 1975 and connection to a deep magma source. *J Volcanol Geotherm Res* 186:379-386, doi: 10.1016/j.jvolgeores.2009.07.006
- Werner C, Hurst T, Scott B, Sherburn S, Christenson BW, Britten K, Cole-Baker J, Mullan B (2008) Variability of passive gas emissions, seismicity, and deformation during crater lake growth at White Island Volcano, New Zealand, 2002–2006. *J Geophys Res* 113:B01204, doi: 10.1029/2007JB005094
- Werner C, Wyngaard JC, Brantley SL (2000) Eddy correlation measurement of hydrothermal gases. *Geophys Res Lett* 27(18):2925-2928, doi: 10.1029/2000GL011765
- Williams SN, Schaefer SJ, Calvache ML, Lopez D (1992) Global carbon dioxide emission to the atmosphere by volcanoes. *Geochim Cosmochim Acta* 56(4):1765-1770, doi: 10.1016/0016-7037(92)90243-C
- Zapata GJA, Calvache VML, Cortés JGP, Fischer TP, Garzon VG, Gómez MD, Narvaez ML, Ordóñez VM, Ortega EA, Stix J, Torres CR, Williams SN (1997) SO₂ fluxes from Galeras volcano, Colombia, 1989-1995: progressive degassing and conduit obstruction of a Decade Volcano. *J Volcanol Geotherm Res* 77:195-208
- Zeebe RE, Caldeira K (2008) Close mass balance of long-term carbon fluxes from ice-core CO₂ and ocean chemistry records. *Nature Geosci* 1:312-315